



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 491/048, A61K 31/415, C07D 491/052, 495/04, 495/14, 513/14 // (C07D 491/048, 307:00, 231:00) (C07D 495/04, 333:00, 231:00) (C07D 491/052, 311:00, 231:00) (C07D 495/04, 335:00, 231:00) (C07D 495/04, 337:00, 231:00) (C07D 495/14, 335:00, 231:00, 221:00)</p>	A1	<p>(11) International Publication Number: WO 96/09304</p> <p>(43) International Publication Date: 28 March 1996 (28.03.96)</p>
<p>(21) International Application Number: PCT/US95/11403</p> <p>(22) International Filing Date: 15 September 1995 (15.09.95)</p> <p>(30) Priority Data: 08/309,291 20 September 1994 (20.09.94) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 08/309,291 (CIP) Filed on 20 September 1994 (20.09.94)</p> <p>(71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): TALLEY, John, J. [US/US]; 8772 Pine Avenue, Brentwood, MO 63144 (US). BERTENSHAW, Stephen, R. [US/US]; 8758 Pine Avenue,</p>		<p>Brentwood, MO 63114 (US). GRANETO, Matthew, J. [US/US]; 1510 Voltaire Drive, St. Louis, MO 63146 (US). ROGIER, Donald, J. [US/US]; 1828 Westmeade Drive, Chesterfield, MO 63017 (US).</p> <p>(74) Agents: BULOCK, Joseph, W. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).</p> <p>(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).</p> <p>Published With international search report.</p>
<p>(54) Title: BENZOPYRANOPYRAZOLYL DERIVATIVES FOR THE TREATMENT OF INFLAMMATION</p> <p>(57) Abstract</p> <p>A class of benzopyranopyrazolyl derivatives is described for use in treating inflammation and inflammation-related disorders. Compounds of particular interest are defined by formula (I) wherein A is $-(CH_2)_m-X-(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from phenyl and five and six membered heteroaryl; wherein R^1 is selected from lower haloalkyl, cyano, lower hydroxyalkyl, formyl, lower alkoxy, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl and lower N-alkyl-N-phenylaminocarbonyl; wherein R^2 is phenyl substituted at a substitutable position with a radical selected from lower alkylsulfonyl and aminosulfonyl; and wherein R^4 is one or more radicals selected from hydrido, halo, lower alkylthio, lower alkylsulfinyl, lower alkyl, cyano, carboxyl, lower alkoxy, aminocarbonyl, lower haloalkyl, hydroxyl, lower alkoxy, amino, lower N-alkylamino, lower N,N-dialkylamino, lower hydroxyalkyl and lower haloalkoxy; or a pharmaceutically-acceptable salt thereof.</p> <div data-bbox="763 877 997 1037"> <p style="text-align: right;">(I)</p> </div>		

Best Available Copy

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**BENZOPYRANOPYRAZOLYL DERIVATIVES
FOR THE TREATMENT OF INFLAMMATION**

FIELD OF THE INVENTION

5

This invention is in the field of antiinflammatory pharmaceutical agents and specifically relates to compounds, compositions and methods for treating inflammation and inflammation-associated disorders, such as arthritis.

10

BACKGROUND OF THE INVENTION

Prostaglandins play a major role in the inflammation process and the inhibition of prostaglandin production, especially production of PGG₂, PGH₂ and PGE₂, has been a common target of antiinflammatory drug discovery. However, common non-steroidal antiinflammatory drugs (NSAIDs) that are active in reducing the prostaglandin-induced pain and swelling associated with the inflammation process are also active in affecting other prostaglandin-regulated processes not associated with the inflammation process. Thus, use of high doses of most common NSAIDs can produce severe side effects, including life threatening ulcers, that limit their therapeutic potential. An alternative to NSAIDs is the use of corticosteroids, which have even more drastic side effects, especially when long term therapy is involved.

Previous NSAIDs have been found to prevent the production of prostaglandins by inhibiting enzymes in the human arachidonic acid/prostaglandin pathway, including the enzyme cyclooxygenase (COX). The recent discovery of an inducible enzyme associated with inflammation (named "cyclooxygenase-2 (COX-2)" or "prostaglandin G/H synthase II") provides a viable target of inhibition which more effectively reduces

35

inflammation and produces fewer and less drastic side effects.

The novel compounds described herein are such safe and also effective antiinflammatory agents. The invention compounds are found to show usefulness in vivo as antiinflammatory agents with minimal side effects. The compounds described herein preferably selectively inhibit cyclooxygenase-2 over cyclooxygenase-1.

Substituted pyrazoles having antiinflammatory activity are described in copending applications 08/160,594 and 08/160,553.

U.S. Patent No. 3,940,418 to R. Hamilton describes tricyclic 4,5-dihydrobenz[g]indazole-3-carboxylic acids as antiinflammatory agents.

U.S. Patent No. 4,803,193 to Kanda et al, describes spiro[3-alkyl-1-aryl[1]benzopyrano[4,3-c]pyrazole-4(1H),9'--[9H]fluorenes as heat sensitive recording materials.

V. Colota et al (*J.Med.Chem.*, 33, 2646 (1991)) describe tricyclic heteroaramatic systems, including 1-aryl-pyrazolo[4,5-c]quinolin-4-ones, 1-aryl-pyrazolo[4,5-c][1,8]naphthyridin-4-ones and 1-aryl-[1]benzopyrano[3,4-d]pyrazol-4-ones for CNS applications. F. Melani et al [*J.Med.Chem.*, 29, 291 (1986)] also describe 1-phenyl-pyrazolo[4,5-c]quinolines for CNS applications.

U.S. Patent Nos. 4,816,467 and 5,206,258 to Doria et al describe (2-cyano-3-(1,4-dihydro)-1-phenyl-[1]benzothiopyrano[4,3-c]pyrazol-3-yl)-3-oxo-propanamides as immunomodulators. G. Doria et al (*Farmaco*, 46, 843 (1991)) also describe the immunomodulating activity of pyrazolylpropanamides, and specifically ethyl [1-(4-fluorophenyl)-1,4-dihydro-[1]benzothiopyrano[4,3-c]pyrazole]-3-carboxylate. British patent 2,227,741 describes related benzopyrano[4,3-c]pyrazoles and benzothiopyrano[4,3-

c]pyrazoles. European application No. 347,773 similarly describes such fused pyrazole compounds, and specifically α -cyano-N,1-bis(4-fluorophenyl)- β -oxo-1H-[1]benzothieno[3,2-c]pyrazole-3-propanamide. U.S.


- 5 Patent No. 5,260,328 to Doria et al describes 2-cyano-3-(1,4-dihydro)-1-phenyl-[1]benzothiopyrano[4,3-c]pyrazol-3-yl)-3-oxo-propanamides for the treatment of rheumatoid arthritis.

- U.S. Patent No. 4,678,499 to Pasteris et al
10 describes 1-aryl-indenopyrazol-4-one-5-sulfonamides as having herbicidal activity. Specifically, 1-phenyl-indenopyrazol-4-one-5-sulfonamide and 1,4-dihydro-N-[[[(4-methoxy-6-methyl-2-pyrimidinyl)amino]carbonyl]-3-methyl-1-[4-(methylsulfonyl)phenyl]-4-oxo-indeno[1,2-
15 c]pyrazole-5-sulfonamide are described.

The invention's benzopyranopyrazolyl derivatives are found to show usefulness *in vivo* as antiinflammatory agents with minimal side effects.

20 DESCRIPTION OF THE INVENTION

A class of compounds useful in treating inflammation-related disorders is defined by Formula I:



(I)

25

- wherein A is $-(CH_2)_m-X-(CH_2)_n-$;
wherein X is selected from $S(O)_p$, O and NR^3 ;
wherein m is 0 to 3, inclusive;
30 wherein n is 0 to 3, inclusive;
wherein p is 0 to 2, inclusive;
wherein B is selected from aryl and heteroaryl;
wherein R^1 is selected from hydrido, halo,
haloalkyl, cyano, nitro, formyl, alkoxycarbonyl,

carboxyl, carboxyalkyl, alkoxycarbonylalkyl, amidino, cyanoamidino, aminocarbonyl, alkoxy, alkoxyalkyl, aminocarbonylalkyl, N-alkylaminocarbonyl, N-arylamino-

5 arylaminocarbonyl, N,N-dialkylaminocarbonyl, N-alkyl-N-arylamino-

10 arylaminocarbonyl, alkylcarbonyl, alkylcarbonylalkyl, hydroxyalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkylthioalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, N-alkylaminosulfonyl, N-arylamino-

15 arylaminosulfonyl, N,N-dialkylaminosulfonyl, N-alkyl-N-arylamino-

20 arylaminosulfonyl and heterocyclic;

wherein R^2 is selected from aryl and heteroaryl, wherein R^2 is optionally substituted at a substitutable position with one or more radicals selected from alkylsulfonyl, aminosulfonyl, halo, alkyl, alkoxy,

15 hydroxyl and haloalkyl;

wherein R^3 is selected from hydrido and alkyl; and wherein R^4 is one or more radicals selected from hydrido, halo, alkylthio, alkylsulfinyl, alkyl, alkylsulfonyl, cyano, carboxyl, alkoxycarbonyl, amido,

20 N-alkylaminocarbonyl, N-arylamino-

25 arylaminocarbonyl, N,N-dialkylaminocarbonyl, N-alkyl-N-arylamino-

30 arylaminocarbonyl, haloalkyl, hydroxyl, alkoxy, hydroxyalkyl, haloalkoxy, aminosulfonyl, N-alkylaminosulfonyl, amino, N-alkylamino, N,N-dialkylamino, heterocyclic, nitro and acylamino;

provided either R^4 is aminosulfonyl or alkylsulfonyl, or R^2 is substituted with aminosulfonyl or alkylsulfonyl;

or a pharmaceutically-acceptable salt thereof.

30 Compounds of Formula I would be useful for, but not limited to, the treatment of inflammation in a subject, and for treatment of other inflammation-associated disorders, such as, as an analgesic in the treatment of pain and headaches, or as an antipyretic

35 for the treatment of fever. For example, compounds of the invention would be useful to treat arthritis, including but not limited to rheumatoid arthritis,

spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus and juvenile arthritis. Such compounds of the invention would be useful in the treatment of asthma, bronchitis, menstrual cramps, tendinitis, bursitis, and skin related conditions such as psoriasis, eczema, burns and dermatitis. Compounds of the invention also would be useful to treat gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome and ulcerative colitis and for the prevention of colorectal cancer. Compounds of the invention would be useful in treating inflammation in such diseases as vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes, myasthenia gravis, multiple sclerosis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, hypersensitivity, conjunctivitis, cystic fibrosis, swelling occurring after injury, myocardial ischemia, and the like. The compounds would also be useful in the treatment of ophthalmic diseases such as retinitis, retinopathies, uveitis, and of acute injury to the eye tissue. The compounds would also be useful for the treatment of certain central nervous system disorders such as alzheimers disease and dementia. The compounds of the invention are useful as anti-inflammatory agents, such as for the treatment of arthritis, with the additional benefit of having significantly less harmful side effects. These compounds would also be useful in the treatment of allergic rhinitis, respiratory distress syndrome, endotoxin shock syndrome, atherosclerosis and central nervous system damage resulting from stroke, ischemia and trauma.

Besides being useful for human treatment, these compounds are also useful for treatment of mammals,

including horses, dogs, cats, rats, mice, sheep, and pigs, and of birds.

- The present compounds may also be used in co-therapies, partially or completely, in place of other conventional antiinflammatories, such as together with
15 steroids, NSAIDs, 5-lipoxygenase inhibitors, LTB₄ antagonists and LTA₄ hydrolase inhibitors.

- Suitable LTB₄ inhibitors include, among others, ebselen, Bayer Bay-x-1005, Ciba Geigy compound CGS-
10 25019C, Leo Denmark compound ETH-615, Lilly compound LY-293111, Ono compound ONO-4057, Terumo compound TMK-688, Lilly compounds LY-213024, 264086 and 292728, ONO compound ONO-LB457, Searle compound SC-53228, calcitrol, Lilly compounds LY-210073, LY223982,
15 LY233469, and LY255283, ONO compound ONO-LB-448, Searle compounds SC-41930, SC-50605 and SC-51146, and SK&F compound SKF-104493. Preferably, the LTB₄ inhibitors are selected from ebselen, Bayer Bay-x-1005, Ciba Geigy compound CGS-25019C, Leo Denmark compound ETH-615,
20 Lilly compound LY-293111, Ono compound ONO-4057, and Terumo compound TMK-688.

- Suitable 5-LO inhibitors include, among others, masoprocol, tenidap, zileuton, pranlukast, tepoxalin, rilopirox, flezelastine hydrochloride, enazadrem
25 phosphate, and bunaprolast.

- The present invention preferably includes compounds which selectively inhibit cyclooxygenase-2 over cyclooxygenase-1. Preferably, the compounds have a cyclooxygenase-2 IC₅₀ of less than about 0.2 μ M, and
30 also have a selectivity ratio of cyclooxygenase-2 inhibition over cyclooxygenase-1 inhibition of at least 50, and more preferably of at least 100. Even more preferably, the compounds have a cyclooxygenase-1 IC₅₀ of greater than about 1 μ M, and more preferably of
35 greater than 10 μ M. Such preferred selectivity may indicate an ability to reduce the incidence of common NSAID-induced side effects.

A preferred class of compounds consists of those compounds of Formula I wherein A is $-(CH_2)_m-X-(CH_2)_n-$; wherein X is selected from $S(O)_p$, O and NR^3 ; wherein m is 0 to 3, inclusive; wherein n is 0 to 3, inclusive; wherein p is 0 to 2, inclusive; wherein B is selected from phenyl, naphthyl and five and six membered heteroaryl; wherein R^1 is selected from halo, lower haloalkyl, cyano, nitro, formyl, lower alkoxy carbonyl, lower carboxyalkyl, lower alkoxy carbonylalkyl, amidino, cyanoamidino, lower alkoxy, lower alkoxyalkyl, aminocarbonyl, lower aminocarbonylalkyl, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl, lower N-alkyl-N-phenylaminocarbonyl, lower alkylcarbonyl, lower alkylcarbonylalkyl, lower hydroxyalkyl, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, lower alkylthioalkyl, lower alkylsulfinylalkyl, lower alkylsulfonylalkyl, lower N-alkylaminosulfonyl, N-phenylaminosulfonyl, phenylsulfonyl, lower N,N-dialkylaminosulfonyl, lower N-alkyl-N-phenylaminosulfonyl and five-seven membered heterocyclic; wherein R^2 is selected from phenyl and five or six membered heteroaryl, wherein R^2 is optionally substituted at a substitutable position with one or more radicals selected from lower alkylsulfonyl, aminosulfonyl, halo, lower alkyl, lower alkoxy, hydroxyl and lower haloalkyl; wherein R^3 is selected from hydrido and lower alkyl; and wherein R^4 is one or more radicals selected from hydrido, halo, lower alkylthio, lower alkylsulfinyl, lower alkyl, lower alkylsulfonyl, cyano, carboxyl, lower alkoxy carbonyl, aminocarbonyl, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl, lower N-alkyl-N-phenylaminocarbonyl, lower haloalkyl, hydroxyl, lower alkoxy, lower hydroxyalkyl, lower haloalkoxy, aminosulfonyl, lower N-alkylaminosulfonyl, amino, lower N-alkylamino, lower N,N-dialkylamino,

five-seven membered heterocyclic, nitro and acylamino; or a pharmaceutically-acceptable salt thereof.

A more preferred class of compounds consists of those compounds of Formula I wherein A is $-(CH_2)_m-X-(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0, 1 or 2; wherein n is 0, 1 or 2; wherein p is 0, 1 or 2; wherein B is selected from phenyl and five and six membered heteroaryl; wherein R^1 is selected from halo, lower haloalkyl, cyano, formyl, lower alkoxy carbonyl, aminocarbonyl, lower alkoxy carbonyl alkyl, lower alkoxy, lower alkoxy alkyl, lower aminocarbonyl alkyl, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl, lower N-alkyl-N-phenylaminocarbonyl and lower hydroxy alkyl; wherein R^2 is phenyl substituted at a substitutable position with a radical selected from lower alkylsulfonyl and aminosulfonyl; and wherein R^4 is one or more radicals selected from hydrido, halo, lower alkylthio, lower alkylsulfinyl, lower alkyl, lower alkylsulfonyl, cyano, carboxyl, lower alkoxy carbonyl, aminocarbonyl, lower N-alkylaminocarbonyl, lower N,N-dialkylaminocarbonyl, lower N-alkyl-N-phenylaminocarbonyl, lower haloalkyl, hydroxyl, lower alkoxy, lower hydroxy alkyl, amino, lower N-alkylamino, lower N,N-dialkylamino, lower haloalkoxy and nitro; or a pharmaceutically-acceptable salt thereof.


An even more preferred class of compounds consists of those compounds of Formula I wherein A is $-(CH_2)_m-X-(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from phenyl and five and six membered heteroaryl; wherein R^1 is selected from lower haloalkyl, lower hydroxy alkyl, cyano, formyl, lower alkoxy carbonyl, lower alkoxy, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl and lower N-alkyl-N-phenylaminocarbonyl; wherein R^2 is phenyl substituted

at a substitutable position with a radical selected from lower alkylsulfonyl and aminosulfonyl; and wherein R^4 is one or more radicals selected from hydrido, halo, lower alkylthio, lower alkylsulfinyl, lower alkyl, cyano, carboxyl, lower alkoxycarbonyl, aminocarbonyl, lower haloalkyl, hydroxyl, lower alkoxy, amino, lower N-alkylamino, lower N,N-dialkylamino, lower hydroxyalkyl and lower haloalkoxy; or a pharmaceutically-acceptable salt thereof.


- 10 A class of compounds of particular interest consists of those compounds of Formula I wherein A is $-(CH_2)_m-X-(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from phenyl, thienyl, pyridyl, furyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, thiaimidazolyl, oxoimidazolyl, azaoxazolyl, azathiazolyl and pyrrolyl; wherein R^1 is selected from fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, hydroxymethyl, hydroxyethyl, cyano, formyl, carboxyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tert-butoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl, aminocarbonyl, methoxy, ethoxy, propoxy, n-butoxy, N-methylaminocarbonyl, N-phenylaminocarbonyl, N,N-dimethylaminocarbonyl, N-methyl-N-phenylaminocarbonyl and methylcarbonyl; wherein R^2 is phenyl substituted at a substitutable position with a radical selected from methylsulfonyl and aminosulfonyl; wherein R^4 is optionally substituted with one or more radicals selected from hydrido, fluoro, chloro, bromo, methylthio, ethylthio, isopropylthio, tert-butylthio,

- isobutylthio, hexylthio, methylsulfinyl, ethylsulfinyl,
 isopropylsulfinyl, *tert*-butylsulfinyl,
 isobutylsulfinyl, hexylsulfinyl, methyl, ethyl,
 isopropyl, *tert*-butyl, isobutyl, hexyl, cyano,
 5 carboxyl, methoxycarbonyl, ethoxycarbonyl,
 isopropoxycarbonyl, *tert*-butoxycarbonyl,
 propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl,
 pentoxycarbonyl, aminocarbonyl, fluoromethyl,
 difluoromethyl, trifluoromethyl, chloromethyl,
 10 dichloromethyl, trichloromethyl, pentafluoroethyl,
 heptafluoropropyl, difluorochloromethyl,
 dichlorofluoromethyl, difluoroethyl, difluoropropyl,
 dichloroethyl, dichloropropyl, hydroxyl, methoxy,
 methylenedioxy, ethoxy, propoxy, *n*-butoxy,
 15 hydroxymethyl and trifluoromethoxy; or a
 pharmaceutically-acceptable salt thereof.


The preferred compounds of Formula I can be
 represented by Formulas Ia-If as follows:




11




(Ig)




(Ih)




(Ii)




(Ij)




(Ik)




(Il)



(Im)




(In)



(Io)

and



(Ip)

10

A family of specific compounds of particular interest within Formula I consists of compounds and pharmaceutically-acceptable salts thereof as shown in the following Tables:

TABLE I

General Structure Ia



5	R ¹	R ²	R ⁴
10	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ H	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ Cl	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ CF ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ H	C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ CH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
15	-CONH ₂	C ₆ H ₅ SO ₂ CH ₃	H
	-CONHCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CONH(C ₆ H ₅)	C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃) ₂	C ₆ H ₅ SO ₂ CH ₃	H
	-CON(C ₂ H ₅) ₂	C ₆ H ₅ SO ₂ CH ₃	H
20	-CON(CH ₃)(C ₂ H ₅)	C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃)(C ₆ H ₅)	C ₆ H ₅ SO ₂ CH ₃	H
		C ₆ H ₅ SO ₂ CH ₃	H
25		C ₆ H ₅ SO ₂ CH ₃	H
	-CN	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OH	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
30	-CH ₂ SOC ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ CH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ CH ₃	C ₆ H ₅ SO ₂ CH ₃	H
35			

TABLE I (cont.)

General Structure Ia

5	R ¹	R ²	R ⁴
	-CH ₂ SO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ C ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F
10	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	7-F
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-F
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	7-Cl
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-Cl
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6,7-(OCH ₂ O)-
15	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-N(CH ₃) ₂
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-Cl, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-Cl, 5-F
20	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5,6-F
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-SCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-SCH ₃
25	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-SOCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-SOCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-CH ₃
	-CF ₃	C ₆ H ₅ F	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ Cl	6-SO ₂ CH ₃
30	-CF ₃	C ₆ H ₅ OCH ₃	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ CH ₃	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ SOCH ₃	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ H	C ₆ H ₅ SO ₂ NH ₂	H
35	-CF ₂ Cl	C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ CF ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ H	C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ CH ₃	C ₆ H ₅ SO ₂ NH ₂	H

TABLE I (cont.)

General Structure Ia



5	R ¹	R ²	R ⁴
	-CO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CONH ₂	C ₆ H ₅ SO ₂ NH ₂	H
	-CONHCH ₃	C ₆ H ₅ SO ₂ NH ₂	H
10	-CONH(C ₆ H ₅)	C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃) ₂	C ₆ H ₅ SO ₂ NH ₂	H
	-CON(C ₂ H ₅) ₂	C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₂ H ₅)	C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₆ H ₅)	C ₆ H ₅ SO ₂ NH ₂	H
15		C ₆ H ₅ SO ₂ NH ₂	H
		C ₆ H ₅ SO ₂ NH ₂	H
	-CN	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OH	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OCH ₃	C ₆ H ₅ SO ₂ NH ₂	H
20	-CH ₂ OC ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SCH ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
25	-CH ₂ SOCH ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ CH ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ C ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
30	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	7-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	7-Cl
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-Cl
35			

TABLE I (cont.)

General Structure Ia

5	R ¹	R ²	R ⁴
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6, 7-(OCH ₂ O)-
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-N(CH ₃) ₂
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-OCH ₃
10	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-Cl, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-Cl, 5-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-CH ₃
15	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5, 6-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-SCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-SCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-SOCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-SOCH ₃
20	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-CH ₃
	-CF ₃	C ₆ H ₅ F	6-SO ₂ NH ₂
	-CF ₃	C ₆ H ₅ Cl	6-SO ₂ NH ₂
	-CF ₃	C ₆ H ₅ OCH ₃	6-SO ₂ NH ₂
	-CF ₃	C ₆ H ₅ CH ₃	6-SO ₂ NH ₂
25	-CF ₃	C ₆ H ₅ SOCH ₃	6-SO ₂ NH ₂
	-CF ₃	thienylSO ₂ NH ₂	5-F, 6-OCH ₃

TABLE II
General Structure Ib

















5	B	R ²
10		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
15		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
20		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂

TABLE II (cont.)










5	B	R^2
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
10		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
15		$-C_6H_5SO_2NH_2$

TABLE III
General Structure Ic



	R ¹	R ²	R ⁴
5	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ H	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ Cl	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ CF ₃	C ₆ H ₅ SO ₂ CH ₃	H
10	-CO ₂ H	C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ CH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CONH ₂	C ₆ H ₅ SO ₂ CH ₃	H
	-CONHCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
15	-CONH(C ₆ H ₅)	C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃) ₂	C ₆ H ₅ SO ₂ CH ₃	H
	-CON(C ₂ H ₅) ₂	C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃)(C ₂ H ₅)	C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃)(C ₆ H ₅)	C ₆ H ₅ SO ₂ CH ₃	H
20		C ₆ H ₅ SO ₂ CH ₃	H
		C ₆ H ₅ SO ₂ CH ₃	H
	-CN	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OH	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
25	-CH ₂ OC ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
30	-CH ₂ SOCH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ CH ₃	C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ CH ₃	H

TABLE III (cont.)
General Structure Ic

	R ¹	R ²	R ⁴
5	-CH ₂ SO ₂ C ₆ H ₅	C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	7-F
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-F
10	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	7-Cl
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-Cl
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6,7-(OCH ₂ O)-
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-N(CH ₃) ₂
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-OCH ₃
15	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-Cl, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-Cl, 5-F
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-CH ₃
20	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5,6-F
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-SCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-SCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	6-SOCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-SOCH ₃
25	-CF ₃	C ₆ H ₅ SO ₂ CH ₃	5-F, 6-CH ₃
	-CF ₃	C ₆ H ₅ F	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ Cl	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ OCH ₃	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ CH ₃	6-SO ₂ CH ₃
30	-CF ₃	C ₆ H ₅ SOCH ₃	6-SO ₂ CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ H	C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ Cl	C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ CF ₃	C ₆ H ₅ SO ₂ NH ₂	H
35	-CO ₂ H	C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ CH ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H

TABLE III (cont.)
General Structure Ic



	R ¹	R ²	R ⁴
5	-CONH ₂	C ₆ H ₅ SO ₂ NH ₂	H
	-CONHCH ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CONH(C ₆ H ₅)	C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃) ₂	C ₆ H ₅ SO ₂ NH ₂	H
10	-CON(C ₂ H ₅) ₂	C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₂ H ₅)	C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₆ H ₅)	C ₆ H ₅ SO ₂ NH ₂	H
		C ₆ H ₅ SO ₂ NH ₂	H
		C ₆ H ₅ SO ₂ NH ₂	H
15	-CN	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OH	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OCH ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
20	-CH ₂ SCH ₃	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ CH ₃	C ₆ H ₅ SO ₂ NH ₂	H
25	-CH ₂ SO ₂ C ₂ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ C ₆ H ₅	C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	7-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-F
30	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	7-Cl
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-Cl
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6,7-(OCH ₂ O)-
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-N(CH ₃) ₂
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-OCH ₃

TABLE III (cont.)
General Structure Ic

5	R ¹	R ²	R ⁴
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-Cl, 6-OCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-Cl, 5-F
10	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-CH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5, 6-F
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-SCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-SCH ₃
15	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	6-SOCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-SOCH ₃
	-CF ₃	C ₆ H ₅ SO ₂ NH ₂	5-F, 6-CH ₃
	-CF ₃	C ₆ H ₅ F	6-SO ₂ NH ₂
	-CF ₃	C ₆ H ₅ Cl	6-SO ₂ NH ₂
20	-CF ₃	C ₆ H ₅ OCH ₃	6-SO ₂ NH ₂
	-CF ₃	C ₆ H ₅ CH ₃	6-SO ₂ NH ₂
	-CF ₃	C ₆ H ₅ SOCH ₃	6-SO ₂ NH ₂
	-CF ₃	thienylSO ₂ NH ₂	5-F, 6-OCH ₃

TABLE IV
General Structure Id















5	B	R^2
10		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
15		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
20		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$

TABLE IV (cont.)










5	B	R^2
10		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
15		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$

TABLE V

General Structure Ie



5	R ¹	R ²	R ⁴
10	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ Cl	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
15	-CONH ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONHCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
20	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
25	-CN	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OH	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
30	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H

TABLE V (cont.)
General Structure Ie

5	R ¹	R ²	R ⁴
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
10	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl
15	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7,8-(OCH ₂ O)-
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-N(CH ₃) ₂
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-OCH ₃
20	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-Cl, 7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl, 6-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6,7-F
25	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
30	-CF ₃	-C ₆ H ₅ F	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ SOCH ₃	7-SO ₂ CH ₃
35			

TABLE V (cont.)

General Structure Ie



5	R ¹	R ²	R ⁴
10	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ Cl	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
15	-CONH ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CONHCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
20		-C ₆ H ₅ SO ₂ NH ₂	H
		-C ₆ H ₅ SO ₂ NH ₂	H
	-CN	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OH	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
25	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
30			

TABLE V (cont.)
General Structure Ia

5	R ¹	R ²	R ⁴
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
10	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl
15	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7,8-(OCH ₂ O)-
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-N(CH ₃) ₂
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-OCH ₃
20	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl, 6-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,7-F
25	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-CH ₃
30	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7,8,9-OCH ₃
	-CF ₃	-C ₆ H ₅ F	7-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ NH ₂
35	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ SOCH ₃	7-SO ₂ NH ₂
	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	7-CH ₃

TABLE VI
General Structure If















5	B	R ²
10		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
15		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
20		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃

TABLE VI (cont.)














	B	R ²
5		
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
10		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
15		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃

TABLE VI (cont.)
















	B	R ²
5		
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
10		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
15		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃

TABLE VI (cont.)












	B	R ²
5		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
10		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
15		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂

TABLE VII
General Structure Ig



5	R ¹	R ²	R ⁴
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ Cl	-C ₆ H ₅ SO ₂ CH ₃	H
10	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONH ₂	-C ₆ H ₅ SO ₂ CH ₃	H
15	-CONHCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
20	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
	-CN	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OH	-C ₆ H ₅ SO ₂ CH ₃	H
25	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
30	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H

TABLE VII (cont.)

General Structure Ig

5	R ¹	R ²	R ⁴
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
10	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl
15	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7, 8- (OCH ₂ O) -
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-N(CH ₃) ₂
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-Cl, 7-OCH ₃
20	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl, 6-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6, 7-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SCH ₃
25	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
	-CF ₃	-C ₆ H ₅ F	7-SO ₂ CH ₃
30	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ SONH ₃	7-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	H
35	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₃	H
	-CF ₂ Cl	-C ₆ H ₅ SO ₂ NH ₃	H
	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ NH ₃	H

TABLE VII (cont.)

General Structure Ig



5	R ¹	R ²	R ⁴
	-CO ₂ H	-C ₆ H ₅ SO ₂ NH ₃	H
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₃	H
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
10	-CONH ₂	-C ₆ H ₅ SO ₂ NH ₃	H
	-CONHCH ₃	-C ₆ H ₅ SO ₂ NH ₃	H
	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₃	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ NH ₃	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ NH ₃	H
15	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ NH ₃	H
	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₃	H
		-C ₆ H ₅ SO ₂ NH ₃	H
		-C ₆ H ₅ SO ₂ NH ₃	H
	-CN	-C ₆ H ₅ SO ₂ NH ₃	H
20	-CH ₂ OH	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ NH ₃	H
25	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
30	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₃	H
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-F

TABLE VII (cont.)

General Structure Ig

5	R ¹	R ²	R ⁴
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	8-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-Cl
10	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	8-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7,8-(OCH ₂ O)-
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-N(CH ₃) ₂
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6-F, 7-OCH ₃
15	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6-Cl, 7-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-Cl, 6-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6-F, 7-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6,7-F
20	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6-F, 7-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6-F, 7-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₃	6-F, 7-CH ₃
25	-CF ₃	-thienylSO ₂ NH ₂	6-F, 7-OCH ₃
	-CF ₃	-C ₆ H ₅ F	7-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ NH ₂
30	-CF ₃	-C ₆ H ₅ SOCH ₃	7-SO ₂ NH ₂

TABLE VIII
General Structure Ih













5	B	R^2
10		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
15		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$

TABLE VIII (cont.)














	B	R ²
5		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
10		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
15		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ CH ₃
		-C ₆ H ₅ SO ₂ NH ₂

TABLE VIII (cont.)
















	B	R ²
5		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
10		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
15		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂

TABLE VIII (cont.)













	B	R ²
5		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
10		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
15		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂

TABLE IX

General Structure Ii



5	P	R ¹	R ²	R ⁴
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CF ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CF ₂ Cl	-C ₆ H ₅ SO ₂ CH ₃	H
10	0	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CO ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CONH ₂	-C ₆ H ₅ SO ₂ CH ₃	H
15	0	-CONHCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
20	0	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
	0		-C ₆ H ₅ SO ₂ CH ₃	H
	0		-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CN	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ OH	-C ₆ H ₅ SO ₂ CH ₃	H
25	0	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
30	0	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
35				

TABLE IX (cont.)

General Structure Ii

5	p	R ¹	R ²	R ⁴
	0	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F
10	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7,8-(OCH ₂ O)-
15	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-N(CH ₃) ₂
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-Cl, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl, 6-F
20	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6,7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SCH ₃
25	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SOCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SOCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
	0	-CF ₃	-C ₆ H ₅ F	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ CH ₃
30	0	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ SOCH ₃	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
35	0	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	7-F
	0	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	7-CH ₃
	0	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	7-Cl
	0	-CF ₂ Cl	-C ₆ H ₅ SO ₂ NH ₂	H

TABLE IX (cont.)

General Structure II



5	P	R ¹	R ²	R ⁴
	0	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CO ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
10	0	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CONH ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CONHCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
15	0	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	0		-C ₆ H ₅ SO ₂ NH ₂	H
	0		-C ₆ H ₅ SO ₂ NH ₂	H
20	0	-CN	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CN	-C ₆ H ₅ SO ₂ NH ₂	7-F
	0	-CN	-C ₆ H ₅ SO ₂ NCHN(CH ₃) ₂	7-F
	0	-CH ₂ OH	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ OH	-C ₆ H ₅ SO ₂ NH ₂	7-F
25	0	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
30	0	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H

TABLE IX (cont.)
General Structure Ii

5	p	R ¹	R ²	R ⁴
	0	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CONH ₂	-C ₆ H ₅ SO ₂ NH ₂	7-F
10	0	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl
15	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-Cl
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7,8-(OCH ₂ O)-
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,7-(OCH ₂ O)-
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-N(CH ₃) ₂
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-OCH ₃
20	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,8-F, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl, 6-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-CH ₃
25	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-SCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-SOCH ₃
30	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SOCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-CH ₃
	0	-CF ₃	thienylSO ₂ NH ₂	6-F, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ F	7-SO ₂ NH ₂
	0	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ NH ₂
35	0	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ NH ₂
	0	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ NH ₂
	0	-CF ₃	-C ₆ H ₅ SOCH ₃	7-SO ₂ NH ₂
	0	-CHF ₂	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-OCH ₃

TABLE IX (cont.)
General Structure Ii

5	p	R ¹	R ²	R ⁴
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-OCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-OCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl, 6-F
10	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-CH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,7-F
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SOCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-CH ₃

TABLE X

General Structure Ij














5	B	R ²	P
10		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
15		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0

TABLE X (cont.)














	B	R ²	p
5		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
10		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
15		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ CH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0

TABLE X (cont.)
















	B	R ²	p
5		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
10		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
15		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0

TABLE X (cont.)
















	B	R ²	p
5		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
10		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
15		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	1
		-C ₆ H ₅ SO ₂ NH ₂	1

TABLE XI

General Structure Ik



5	P	R ¹	R ²	R ⁴
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CF ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CF ₂ Cl	-C ₆ H ₅ SO ₂ CH ₃	H
10	0	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CO ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CONH ₂	-C ₆ H ₅ SO ₂ CH ₃	H
15	0	-CONHCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CONH(C ₆ H ₃)	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
20	0	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
	0		-C ₆ H ₅ SO ₂ CH ₃	H
	0		-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CN	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ OH	-C ₆ H ₅ SO ₂ CH ₃	H
25	0	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
30	0	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
35	0	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H

TABLE XI (cont.)

General Structure Ik

5	p	R ¹	R ²	R ⁴
	0	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F
10	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7,8-(OCH ₂ O)-
15	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-N(CH ₃) ₂
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-Cl, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl, 6-F
20	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6,7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SCH ₃
25	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-SOCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-SOCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	6-F, 7-CH ₃
	0	-CF ₃	-C ₆ H ₅ F	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ CH ₃
30	0	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ SOCH ₃	7-SO ₂ CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
35	0	-CF ₂ Cl	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	CO ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H

TABLE XI (cont.)

General Structure Ik



5	p	R ¹	R ²	R ⁴
	0	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CONH ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CONHCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
10	0	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
15	0		-C ₆ H ₅ SO ₂ NH ₂	H
	0		-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CN	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ OH	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
20	0	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
25	0	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
30	0	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl
35				

TABLE XI (cont.)

General Structure Ik

5	p	R ¹	R ²	R ⁴
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-Cl
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7,8-(OCH ₂ O)-
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-N(CH ₃) ₂
10	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7,8-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl, 6-F
15	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-CH(CH ₃) ₂
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-CH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,7-F
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,7-Cl
20	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-SCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-SOCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SOCH ₃
	0	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-CH ₃
25	0	-CF ₃	-thienylSO ₂ NH ₂	6-F, 7-OCH ₃
	0	-CF ₃	-C ₆ H ₅ F	7-SO ₂ NH ₂
	0	-CF ₃	-C ₆ H ₅ Cl	7-SO ₂ NH ₂
	0	-CF ₃	-C ₆ H ₅ OCH ₃	7-SO ₂ NH ₂
	0	-CF ₃	-C ₆ H ₅ CH ₃	7-SO ₂ NH ₂
30	0	-CF ₃	-C ₆ H ₅ SOCH ₃	7-SO ₂ NH ₂
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-OCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-OCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl, 6-F
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-CH ₃
35	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6,7-F
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-F, 7-SOCH ₃
	1	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	6-Cl, 7-CH ₃

TABLE XII
General Structure I1













5	B	R ²	p
10		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
15		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0

TABLE XII (cont.)














	B	R ²	P
5		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
10		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
15		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ CH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0

TABLE XII (cont.)















	B	R ²	p
5		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
10		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
15		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0

TABLE XII (cont.)
















	B	R ²	p
5		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
10		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₃	0
15		-C ₆ H ₅ SO ₂ NH ₃	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0
		-C ₆ H ₅ SO ₂ NH ₂	0

TABLE XIII

General Structure Im



5	R ¹	R ²	R ⁴
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ Cl	-C ₆ H ₅ SO ₂ CH ₃	H
10	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONH ₂	-C ₆ H ₅ SO ₂ CH ₃	H
15	-CONHCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONH(C ₆ H ₃)	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
20	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
	-CN	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OH	-C ₆ H ₅ SO ₂ CH ₃	H
25	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
30	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H

TABLE XIII (cont.)

General Structure Im

5	R ¹	R ²	R ⁴
	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
10	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	9-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	9-Cl
15	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8,9-(OCH ₂ O)-
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-N(CH ₃) ₂
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl, 8-OCH ₃
20	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl, 7-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7,8-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-SCH ₃
25	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-CH ₃
	-CF ₃	-C ₆ H ₅ F	8-SO ₂ CH ₃
30	-CF ₃	-C ₆ H ₅ Cl	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ OCH ₃	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ CH ₃	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ SOCH ₃	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
35	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ Cl	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H

TABLE XIII (cont.)

General Structure Im



5	R ¹	R ²	R ⁴
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
10	-CONH ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CONHCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CONH(C ₆ H ₃)	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
15	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
		-C ₆ H ₅ SO ₂ NH ₂	H
		-C ₆ H ₅ SO ₂ NH ₂	H
	-CN	-C ₆ H ₅ SO ₂ NH ₂	H
20	-CH ₂ OH	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
25	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
30	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F
35	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-F

TABLE XIII (cont.)

General Structure Im

5	R ¹	R ²	R ⁴
	-GF ₃	-C ₆ H ₅ SO ₂ NH ₂	9-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7,8-(OCH ₂ O)-
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-N(CH ₃) ₂
10	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl, 8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-Cl, 7-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-CH ₃
15	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7,8-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-SOCH ₃
20	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-CH ₃
	-CF ₃	thienylSO ₂ NH ₂	7-F, 9OCH ₃
	-CF ₃	-C ₆ H ₅ F	8-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ Cl	8-SO ₂ NH ₂
25	-CF ₃	-C ₆ H ₅ OCH ₃	8-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ CH ₃	8-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ SOCH ₃	8-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	9-Cl
30			

TABLE XIV
General Structure In
















5	B	R^2
10		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
15		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
20		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2NH_2$

TABLE XIV (cont.)












	B	R ²
5		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
10		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂
15		-C ₆ H ₅ SO ₂ NH ₂
		-C ₆ H ₅ SO ₂ NH ₂

TABLE XV

General Structure Io



5	R ¹	R ²	R ⁴
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₂ Cl	-C ₆ H ₅ SO ₂ CH ₃	H
10	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ H	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONH ₂	-C ₆ H ₅ SO ₂ CH ₃	H
15	-CONHCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CONH(C ₆ H ₃)	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ CH ₃	H
	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
20	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
		-C ₆ H ₅ SO ₂ CH ₃	H
	-CN	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OH	-C ₆ H ₅ SO ₂ CH ₃	H
25	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
30	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ CH ₃	H

TABLE XV (cont.)
General Structure Io

5	R ¹	R ²	R ⁴
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ CH ₃	H
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F
10	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	9-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	9-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8,9-(OCH ₂ O)-
15	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-N(CH ₃) ₂
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-Cl, 8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-Cl, 7-F
20	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7,8-F
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-SCH ₃
25	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	8-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ CH ₃	7-F, 8-CH ₃
	-CF ₃	-C ₆ H ₅ F	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ Cl	8-SO ₂ CH ₃
30	-CF ₃	-C ₆ H ₅ OCH ₃	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ CH ₃	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ SOCH ₃	8-SO ₂ CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
35	-CF ₂ Cl	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₂ CF ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ H	-C ₆ H ₅ SO ₂ NH ₂	H
	-CO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H

TABLE XV (cont.)
General Structure Io



5	R ¹	R ²	R ⁴
	-CO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CONH ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CONHCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
10	-CONH(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(C ₂ H ₅) ₂	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₂ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
	-CON(CH ₃)(C ₆ H ₅)	-C ₆ H ₅ SO ₂ NH ₂	H
			
15		-C ₆ H ₅ SO ₂ NH ₂	H
			
		-C ₆ H ₅ SO ₂ NH ₂	H
	-CN	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OH	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
20	-CH ₂ OC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ OC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
25	-CH ₂ SOCH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SOC ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ CH ₃	-C ₆ H ₅ SO ₂ NH ₂	H
	-CH ₂ SO ₂ C ₂ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
30	-CH ₂ SO ₂ C ₆ H ₅	-C ₆ H ₅ SO ₂ NH ₂	H
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	9-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-Cl

TABLE XV (cont.)
General Structure Io

5	R ¹	R ²	R ⁴
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	9-Cl
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7, 8-(OCH ₂ O)-
10	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-N(CH ₃) ₂
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 9-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-Cl, 8-OCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-Cl, 7-F
15	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-CH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7, 8-F
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-SCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-SCH ₃
20	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	8-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-SOCH ₃
	-CF ₃	-C ₆ H ₅ SO ₂ NH ₂	7-F, 8-CH ₃
	-CF ₃	thienylSO ₂ NH ₂	7-F, 8-OCH ₃
	-CF ₃	-C ₆ H ₅ F	8-SO ₂ NH ₂
25	-CF ₃	-C ₆ H ₅ Cl	8-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ OCH ₃	8-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ CH ₃	8-SO ₂ NH ₂
	-CF ₃	-C ₆ H ₅ SOCH ₃	8-SO ₂ NH ₂

TABLE XVI
General Structure Ip
























5	B	R^2
10		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
15		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2CH_3$
		$-C_6H_5SO_2NH_2$

TABLE XVI (cont.)

5	B	R^2
10		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
		$-C_6H_5SO_2NH_2$
15		$-C_6H_5SO_2NH_2$

Within Formula I there is a subclass of compounds of high interest represented by Formula II:



5

wherein A is $-(CH_2)_m-O-(CH_2)_n-$ or $-(CH_2)_m-S(O)_p-(CH_2)_n-$; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from aryl and heteroaryl; wherein R^1 is selected from haloalkyl, hydroxyalkyl, aminocarbonyl, alkoxycarbonyl and cyano; wherein R^4 is one or more radicals selected from hydrido, halo, alkyl and alkoxy; and wherein R^5 is selected from alkyl and amino; or a pharmaceutically-acceptable salt thereof.

A preferred class of compounds consists of those compounds of Formula II wherein A is $-(CH_2)_m-O-(CH_2)_n-$ or $-(CH_2)_m-S(O)_p-(CH_2)_n-$; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from phenyl and five membered heteroaryl; wherein R^1 is selected from lower haloalkyl, lower hydroxyalkyl, aminocarbonyl, lower alkoxycarbonyl and cyano; wherein R^4 is one or more radicals selected from hydrido, halo, lower alkyl and lower alkoxy; and wherein R^5 is selected from lower alkyl and amino; or a pharmaceutically-acceptable salt thereof.

A class of compounds of particular interest consists of those compounds of Formula II wherein A is $-(CH_2)_m-O-(CH_2)_n-$ or $-(CH_2)_m-S(O)_p-(CH_2)_n-$; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1;

30

wherein B is selected from phenyl, thienyl, furyl and pyrrolyl; wherein R¹ is selected from fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, hydroxymethyl, hydroxyethyl, aminocarbonyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and cyano; wherein R⁴ is one or more radicals selected from hydrido, fluoro, chloro, bromo, methyl, ethyl, isopropyl, tert-butyl, isobutyl, hexyl, methoxy, methylenedioxy, ethoxy, propoxy, n-butoxy and tert-butoxy; and wherein R⁵ is selected from methyl and amino; or a pharmaceutically-acceptable salt thereof.

A family of specific compounds of particular interest within Formula II consists of compounds and pharmaceutically-acceptable salts thereof as follows:

- 20 4-[3-(difluoromethyl)-1,5-dihydro-7-methyl-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
7-fluoro-1,5-dihydro-1-[4-(methylsulfonyl)phenyl]-3-
25 (trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazole;
4-[1,5-dihydro-7,8,9-trimethoxy-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
4-[6,8-difluoro-1,5-dihydro-7-methoxy-3-
30 (trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-
1-yl]benzenesulfonamide;
4-[3-cyano-7-fluoro-1,5-dihydro-[2]benzothiopyrano[4,3-
c]pyrazol-1-yl]benzenesulfonamide;
4-[3-(difluoromethyl)-7-fluoro-1,5-dihydro-
35 [2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;

- 4-[1,5-dihydro-7-methyl-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[3-cyano-7-fluoro-1,5-dihydro-[2]benzothiopyrano[4,3-
c]pyrazol-1-yl]-N-
[(dimethylamino)methylene]benzenesulfonamide;
- 4-[3-(difluoromethyl)-1,5-dihydro-7-methyl-
[2]benzenethiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 10 4-[7-fluoro-1,5-dihydro-3-(hydroxymethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,5-dihydro-3-(trifluoromethyl)-
[1,3]dioxolo[6,7][2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 15 4-[1,5-dihydro-7-methoxy-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[7-chloro-3-(difluoromethyl)-1,5-dihydro-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 20 4-[7-chloro-1,5-dihydro-3-trifluoromethyl-
thieno[3',2':4,5]thiopyrano-s-oxide[3,2-c]pyrazol-
1-yl]benzenesulfonamide;
- 25 methyl [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-
fluoro-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-
c]pyrazol-3-yl]carboxylate;
- [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-fluoro-3-
(trifluoromethyl)-[2]benzothiopyrano[4,3-
c]pyrazol-3-yl]carboxamide;
- 30 4-[1,5-dihydro-6-fluoro-7-methoxy-3-
(difluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-
1-yl]benzenesulfonamide;
- 4-[1,5-dihydro-7-fluoro-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 35

- 4-[1,5-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 1,5-dihydro-6-fluoro-7-methoxy-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazole;
- 4-[1,4-dihydro-3-(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 10 methyl [1-[4-(aminosulfonyl)phenyl]-1,4-dihydro-[1]benzopyrano[4,3-c]pyrazol-3-yl]carboxylate;
- 4-[1,4-dihydro-6-fluoro-3-(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 15 4-[3-(trifluoromethyl)-1H-benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,4-dihydro-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 20 methyl [1-[4-(aminosulfonyl)phenyl]-1,4-dihydro-[1]benzothiopyrano[4,3-c]pyrazol-3-carboxylate;
- 4-[6,7-dichloro-1,4-dihydro-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 25 4-[1,4-dihydro-7-fluoro-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,4-dihydro-6-isopropyl-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 30 4-[1,4-dihydro-7,8-dimethoxy-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,4-dihydro-7-methoxy-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 35

- 4-[1,4-dihydro-7-methyl-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 5 4-[7-chloro-1,4-dihydro-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,5-dihydro-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 10 4-[1,5-dihydro-7-methyl-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 1,5-dihydro-1-[4-(methylsulfonyl)phenyl]-7-methyl-3-
(trifluoromethyl)-[2]benzothiopyrano[4,3-
15 c]pyrazole;
- 4-[7-chloro-1,5-dihydro-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,5-dihydro-7-methoxy-3-(trifluoromethyl)-
20 [2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[7-chloro-1,5-dihydro-3-trifluoromethyl-
[2]thienothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide; and
- 25 4-[3-cyano-1,4-dihydro[1]benzothiopyrano[4,3-
c]pyrazol-1-yl]benzenesulfonamide.

The term "hydrido" denotes a single hydrogen atom (H). This hydrido radical may be attached, for example,
30 to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH₂-) radical. Where used, either alone or within other terms such as "haloalkyl", "alkylsulfonyl", "alkoxyalkyl" and "hydroxyalkyl", the
35 term "alkyl" embraces linear or branched radicals having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about

ten carbon atoms. Most preferred are lower alkyl radicals having one to about six carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl and the like. The term "halo" means halogens such as fluorine, chlorine, bromine or iodine. The term "haloalkyl" embraces radicals wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically embraced are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalo and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. "Lower haloalkyl" embraces radicals having 1-6 carbon atoms. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl. The term "hydroxyalkyl" embraces linear or branched alkyl radicals having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms and one or more hydroxyl radicals. Examples of such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl. The terms "alkoxy" and "alkoxyalkyl" embrace linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals are "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and tert-butoxy. The term "alkoxyalkyl" embraces alkyl radicals having one or more alkoxy

radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals. More preferred alkoxyalkyl radicals are "lower alkoxyalkyl" radicals having one to six carbon atoms and one or two alkoxy radicals. Examples of such radicals include methoxymethyl, methoxyethyl, ethoxyethyl, methoxybutyl and methoxypropyl. The "alkoxy" or "alkoxyalkyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide "haloalkoxy" or haloalkoxyalkyl radicals. More preferred haloalkoxy radicals are "lower haloalkoxy" radicals having one to six carbon atoms and one or more halo radicals. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy. The term "aryl", alone or in combination, means a carbocyclic aromatic system containing one, two or three rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" embraces aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane and biphenyl. The term "heterocyclic" embraces saturated, partially saturated and unsaturated heteroatom-containing ring-shaped radicals, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Such heterocyclic radicals preferably include ring systems having 3 to 10 members. Examples of saturated heterocyclic radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms [e.g. pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl, etc.]; saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. morpholinyl, etc.]; saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., thiazolidinyl, etc.]. Examples of partially saturated heterocyclic radicals include dihydrothiophene, dihydropyran, dihydrofuran and dihydrothiazole.

Examples of unsaturated heterocyclic radicals, also termed "heteroaryl" radicals, include unsaturated 3 to 6 membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazolyl [e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.] tetrazolyl [e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.], etc.; unsaturated condensed heterocyclic group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indoliziny, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl [e.g., tetrazolo[1,5-b]pyridazinyl, etc.], etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.; unsaturated 3 to 6-membered heteromonocyclic group containing a sulfur atom, for example, thienyl, etc.; unsaturated 3- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl [e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.] etc.; unsaturated condensed heterocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms [e.g. benzoxazolyl, benzoxadiazolyl, etc.]; unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl [e.g., 1,2,4- thiadiazolyl, 1,3,4- thiadiazolyl, 1,2,5-thiadiazolyl, etc.] and isothiazolyl; unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms [e.g., benzothiazolyl, benzothiadiazolyl, etc.] and the like. The term also embraces radicals where heterocyclic radicals are fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuryl, benzothieryl, and the like. Said "heterocyclic" radicals may have 1 to 3 substituents such as lower alkyl, hydroxy, oxo, amino and lower

alkylamino. More preferred heteroaryl radicals include five to six membered heteroaryl radicals. The term "alkylthio" embraces radicals containing a linear or branched alkyl radical, of one to about ten carbon atoms attached to a divalent sulfur atom. More preferred alkylthio radicals are "lower alkylthio" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthio radicals are methylthio, ethylthio, propylthio, butylthio and hexylthio. The term "alkylthioalkyl" embraces alkylthio radicals attached to an alkyl radical. More preferred alkylthioalkyl radicals are "lower alkylthioalkyl" radicals having alkyl radicals of one to six carbon atoms and an alkylthio radical as described above. Examples of such radicals include methylthiomethyl. The term "arylthio" embraces radicals containing an aryl radical, attached to a divalent sulfur atom, such as a phenylthio radical. The term "alkylsulfinyl" embraces radicals containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent $-S(=O)-$ radical. More preferred alkylsulfinyl radicals are "lower alkylsulfinyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfinyl radicals include methylsulfinyl, ethylsulfinyl, butylsulfinyl and hexylsulfinyl. The term "alkylsulfinylalkyl" embraces alkylsulfinyl radicals attached to an alkyl radical, where alkyl and alkylsulfinyl are defined as above. More preferred alkylsulfinylalkyl radicals are "lower alkylsulfinylalkyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfinylalkyl radicals include methylsulfinylmethyl. The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, denotes respectively divalent radicals $-SO_2-$. "Alkylsulfonyl" embraces alkyl radicals attached to a sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon

- atoms. Examples of such lower alkylsulfonyl radicals include methylsulfonyl, ethylsulfonyl and propylsulfonyl. The term "alkylsulfonylalkyl" embraces alkylsulfonyl radicals attached to an alkyl radical,
- 5 where alkyl and alkylsulfonyl are defined as above. More preferred alkylsulfonylalkyl radicals are "lower alkylsulfonylalkyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfonylalkyl radicals include methylsulfonylmethyl,
- 10 ethylsulfonylmethyl and propylsulfonylmethyl. The term "arylsulfonyl" embraces aryl radicals as defined above, attached to a sulfonyl radical. Examples of such radicals include phenylsulfonyl. The terms "sulfamyl", "aminosulfonyl" and "sulfonamidyl" denotes $\text{NH}_2\text{O}_2\text{S}-$.
- 15 The terms "N-alkylaminosulfonyl" and "N,N-dialkylaminosulfonyl" denote aminosulfonyl radicals substituted, respectively, with one alkyl radical, a cycloalkyl ring, or two alkyl radicals. The terms "N-arylamino sulfonyl" and "N-alkyl-N-arylamino sulfonyl" denote aminosulfonyl radicals substituted with one aryl radical or one alkyl and one aryl radical,
- 20 respectively. The term "acyl" denotes a radical provided by the residue after removal of hydroxyl from an organic acid. Examples of such acyl radicals include
- 25 formyl, alkanoyl and aroyl radicals. The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", denotes $-\text{CO}_2\text{H}$. The term "carbonyl", whether used alone or with other terms, such as "alkoxycarbonyl", denotes $-(\text{C}=\text{O})-$. The
- 30 term "alkoxycarbonyl" means a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl radical. Preferably, "lower alkoxycarbonyl" embraces alkoxy radicals having one to six carbon atoms. Examples of such "lower
- 35 alkoxycarbonyl" ester radicals include substituted or unsubstituted methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and hexyloxycarbonyl. The term "alkylcarbonyl" includes radicals having

alkyl, aryl and aralkyl radicals, respectively, as defined above, attached via an oxygen atom to a carbonyl radical. More preferred alkylcarbonyl radicals are "lower alkylcarbonyl" radicals having one to six carbon atoms. Examples of such radicals include methylcarbonyl and ethylcarbonyl. The term "alkylcarbonylalkyl" embraces radicals having "alkylcarbonyl", as defined above substituted to an alkyl radical. More preferred alkylcarbonylalkyl radicals are "lower alkylcarbonylalkyl" having lower alkylcarbonyl radicals as defined above attached to one to six carbon atoms. Examples of such lower alkylcarbonylalkyl radicals include methylcarbonylmethyl. The term "alkoxycarbonylalkyl" embraces radicals having "alkoxycarbonyl", as defined above substituted to an alkyl radical. More preferred alkoxycarbonylalkyl radicals are "lower alkoxycarbonylalkyl" having lower alkoxycarbonyl radicals as defined above attached to one to six carbon atoms. Examples of such lower alkoxycarbonylalkyl radicals include methoxycarbonylmethyl. The term "carboxyalkyl" embraces radicals having a carboxy radical as defined above, attached to an alkyl radical. More preferred are "lower carboxyalkyl" having alkyl portions of one to six carbon atoms. The term "aminoalkyl" embraces alkyl radicals substituted with amino radicals. More preferred aminoalkyl radicals are "lower aminoalkyl" having one to six carbon atoms. Examples include aminomethyl, aminoethyl and aminobutyl. The term "alkylamino" denotes amino groups which have been substituted with one or two alkyl radicals. Preferred alkylamino radicals are "lower alkylamino" having alkyl portions of one to six carbon atoms. Examples include N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino and the like. The term "alkylaminocarbonyl" embraces alkylamino radicals, as described above, attached to a carbonyl radical. More preferred alkylaminocarbonyl radicals are "lower

alkylaminocarbonyl" having lower alkylamino radicals, as described above, attached to a carbonyl radical. Examples of such radicals include N-methylaminocarbonyl and N,N-dimethylcarbonyl. The term "arylamino" denotes amino groups which have been substituted with one or two aryl radicals, such as N-phenylamino. The "arylamino" radicals may be further substituted on the aryl ring portion of the radical. The term "arylaminoalkyl" embraces arylamino radicals, as described above, connected to a carbonyl radical. An example of such radicals includes phenylaminocarbonyl. The term "N-alkyl-N-arylaminoalkyl" embraces arylamino radicals, as described above, to a carbonyl radical. An example of such radicals includes phenylaminocarbonyl. The term "aminocarbonyl" denotes an amide group of the formula $-C(=O)NH_2$. The term "N-alkyl-N-arylaminoalkyl" embraces aminocarbonyl radicals, as described above, substituted with one alkyl and one aryl radical. An example of such radicals is N-methyl-N-phenylaminocarbonyl. The term "aminocarbonylalkyl" denotes an aminocarbonyl radical attached to an alkyl radical, as described above. An example of such radicals is aminocarbonylmethyl. The term "amidino" denotes an $-C(=NH)-NH_2$ radical. The term "cyanoamidino" denotes an $-C(=N-CN)-NH_2$ radical. The term "cycloalkyl" embraces radicals having three to ten carbon atoms, such as cyclopropyl cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The term "acylamino" embraces an amino radical substituted with an acyl group. An examples of an "acylamino" radical is acetyl amino ($CH_3C(=O)-NH-$).

The present invention comprises a pharmaceutical composition comprising a therapeutically-effective amount of a compound of Formula I in association with at least one pharmaceutically-acceptable carrier, adjuvant or diluent.

The present invention also comprises a method of treating inflammation or inflammation-associated disorders in a subject, the method comprising administering to the subject having such

- 5 inflammation or disorder a therapeutically-effective amount of a compound of Formula I.

Compounds of Formula I would also be capable of inhibiting cytokines, such as TNF, IL-1, IL-6, and IL-8. As such, the compounds can be used in the manufacture of
10 a medicament or in a method for the treatment for the prophylactic or therapeutic treatment of diseases mediated by cytokines, such as TNF, IL-1, IL-6, and IL-8.


- Also included in the family of compounds of
15 Formula I are the pharmaceutically-acceptable salts thereof. The term "pharmaceutically-acceptable salts" embraces salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt is not critical, provided that
20 it is pharmaceutically-acceptable. Suitable pharmaceutically-acceptable acid addition salts of compounds of Formula I may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic,
25 hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, example of which are formic, acetic,
30 propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, salicylic, p-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic),
35 methanesulfonic, ethylsulfonic, benzenesulfonic, pantothenic, toluenesulfonic, 2-hydroxyethanesulfonic, sulfanilic, stearic, cyclohexylaminosulfonic, algenic, β -hydroxybutyric, salicylic, galactaric and

galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of Formula I include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, choline, chloroprocaine, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound of Formula I by reacting, for example, the appropriate acid or base with the compound of Formula I.

GENERAL SYNTHETIC PROCEDURES


The compounds of the invention can be synthesized according to the following procedures of Schemes I-XII, wherein the R¹-R⁵ substituents are as defined for Formulas I-II, above, except where further noted.

SCHEME I



Synthetic Scheme I illustrates the procedure used to prepare the antiinflammatory pyrazoles 3 of the present invention. 1,3-Dicarbonyl compounds such as 1, or the shown enol form which is in equilibrium to the diketone, are reacted with a hydrazine hydrochloride 2 in warm ethanol to provide the pyrazoles 3 via a condensation reaction.

SCHEME II




5 Synthetic Scheme II illustrates the four step
procedure for the preparation of substituted diketones
8. In step one, an appropriately substituted methyl
halide 4 (where X is chloro for example) is converted
into the corresponding thiouronium salt 5 upon
10 treatment with thiourea. In step two, the thiouronium
salt 5 is converted according to the procedure of Lumma
and Berchtold (*J. Org. Chem.*, 34, 1566 (1969)) to the
free mercaptide and then trapped with chloroacetic acid
or a related salt to provide the acetic acid

derivatives 6. In step three, the acids 6 are reacted with trifluoroacetic anhydride (TFAA) in trifluoroacetic acid (TFA) to give the ketones 7. In step four, the ketones 7 are first treated with base, such as sodium methoxide or lithium diisopropylamide (LDA), followed by condensation with a suitable acylating agent, R^1COLG , (where LG represents an appropriate leaving group such as methoxy, chloro, imidazole and the like) in an appropriate solvent, such as methanol, diethyl ether or tetrahydrofuran, to provide the desired diketones 8 which are suitable for conversion into antiinflammatory pyrazoles as illustrated in Scheme I.

15

SCHEME III




Synthetic Scheme III illustrates the four step procedure for the preparation of substituted isothiochromanone 1,3-carbonyl derivatives 13. In step one, an appropriately substituted benzyl alcohol 9 is converted into the corresponding benzyl chloride by stirring with concentrated hydrochloric acid and then

immediately converted into a thiouronium salt 10 upon treatment with thiourea at reflux. In step two, the thiouronium salt is converted to the free mercaptide and then trapped with chloroacetic acid or a related salt to provide the acetic acid derivatives 11. In step three, the acids 11 are reacted with trifluoroacetic anhydride (TFAA) in trifluoroacetic acid (TFA) to give the isothiochromanone products 12. In step four, the isothiochromanones 12 are first treated with base, such as sodium methoxide, sodium bistrimethylsilylamide or lithium diisopropylamide (LDA), followed by condensation with a suitable acylating agent, R^1COLG , (where LG is defined as in Scheme II) in an appropriate solvent, such as methanol, diethyl ether or tetrahydrofuran, to provide 1,3-dicarbonyl compounds 13 which are suitable for conversion into antiinflammatory pyrazoles as illustrated in Scheme I.

Alternatively, the dicarbonyl compounds 13 can be directly prepared from commercially available isothiochromanones 12. The thiouronium salts 10 can be prepared from commercially available benzyl halides.

SCHEME IV




Synthetic Scheme IV illustrates a three step

procedure used for the preparation of substituted thiochromanone 1,3-dicarbonyl derivatives 17. In step one, an appropriate substituted thiophenol 14 is converted into the corresponding propionic acid derivatives 15 upon treatment with acrylic acid at a temperature in a range of room temperature to about 50°C. In step two, the propionic acids 15 are subjected to treatment with a mixture of trifluoroacetic anhydride and trifluoroacetic acid to effect intramolecular Friedel-Crafts acylation, thus providing thiochromanones 16. In the last step, substituted thiochromanones 16 are first treated with a base, such as lithium diisopropyl amide or sodium methoxide (LDA), followed by condensation with suitable acylating agents $R^1\text{COLG}$ (as defined in Scheme II) in an appropriate solvent such as diethylether, methanol or tetrahydrofuran to provide the 1,3-dicarbonyl compounds 17 which are suitable for conversion into antiinflammatory pyrazoles as illustrated in Scheme I.

Alternatively, the dicarbonyl compounds 17 can be directly prepared from commercially available thio-4-chromanones 16.


SCHEME V



Synthetic Scheme V details the three step procedure used to prepare substituted 1,3-dicarbonyl chromanone derivatives 21. In step one, substituted phenols 18 are condensed with acrylic acid to afford 3-phenoxypropionic acids 19. In step two, the acids 19 are treated with a mixture of trifluoroacetic anhydride and trifluoroacetic acid to affect intramolecular Friedel-Crafts acylation affording selected chromanones 20. In step three, substituted chromanones 20 are first treated with base, such as lithium diisopropylamide (LDA) or sodium methoxide followed by condensation with suitable acylating agents, $R^1\text{COLG}$ (where LG represents leaving group as previously defined in Scheme II in an appropriate solvent such as diethyl ether or methanol) to provide 1,3-dicarbonyl compounds 21 which are suitable for conversion into antiinflammatory pyrazoles as illustrated in Scheme I.

Alternatively, the dicarbonyl compounds 21 can be directly formed from commercially available chromanones 20.


SCHEME VI



Synthetic Scheme VI illustrates a three step procedure used to prepare substituted 1,3-dicarbonyl

isochromanone derivatives 25. In step one, selected benzyl alcohol derivatives 22 are treated with sodium hydride and subsequently treated with ethyl bromoacetate to provide the desired ethers 23. In step two, the ester group of 23 is hydrolyzed with aqueous sodium hydroxide and then treated with a mixture of trifluoroacetic acid and trifluoroacetic anhydride to promote intramolecular Friedel-Crafts acylation affording isochromanone 24 derivatives. In the third step, the isochromanones 24 are first treated with a base such as lithium diisopropylamide (LDA) or sodium methoxide followed by condensation with suitable acylating agents (R^1COLG) to provide the 1,3-dicarbonyl compounds 25 which were suitable for conversion into antiinflammatory pyrazoles as illustrated in Scheme I.


SCHEME VII



20

Synthetic Scheme VII illustrates a procedure used to prepare the methylsulfonylphenylhydrazine hydrochloride and the sulfonamidyphenylhydrazine hydrochlorides 27 as used in Scheme I. The sulfonylphenylhydrazine 26 is converted to the hydrochloride salt by stirring with a 4N solution of hydrochloric acid in a solvent such as dioxane.


SCHEME VIII



5 Synthetic Scheme VIII illustrates a procedure used
to prepare substituted 3-coumaranones 29. Coumaranones
28 are first treated with a base, such as lithium
diisopropyl amide or sodium methoxide (LDA) followed by
condensation with suitable acylating agents R¹COLG (as
10 defined in Scheme II) in an appropriate solvent such as
diethylether, methanol or tetrahydrofuran to provide
the 1,3-dicarbonyl compounds 29 which are suitable for
conversion into antiinflammatory pyrazoles as
illustrated in Scheme I.

15

SCHEME IX




20 Synthetic Scheme IX illustrates a two step
procedure used for the preparation of substituted
benzylalcohols 9. In step one, a mixture of potassium

tert-butoxide and anhydrous tetrahydrofuran, cooled to -78°C and treated with a 1.6 M solution of n-butyllithium in hexanes, is added to an appropriate substituted benzene 30 the anion thereby generated is
 5 reacted with carbon dioxide to yield the benzoic acid 31. In step two, the benzoic acid 31 is dissolved in a solvent, such as tetrahydrofuran, and treated with a reducing agent, such as borane dimethyl sulfide complex, to form the desired benzyl alcohol 9.


10

SCHEME X




15 Synthetic Scheme X illustrates a procedure used for the preparation of the antiinflammatory oxidized thio-containing fused tricyclic pyrazoles 33. The appropriate pyrazole 32 from Scheme I, where A is S or -(CH₂)_mS(CH₂)_n-, is treated with an oxidizing agent
 20 such as *m*-chloroperbenzoic acid (MCPBA) or hydrogen peroxide. Compounds having differing amounts of oxidation (sulfinyls and sulfones) can be separated, such as by chromatography.

SCHEME XI




- 5 Synthetic Scheme XI shows procedures for preparing antiinflammatory agents 36, 37 and 38 of Formula I. The esters 35, which can be prepared as shown in Scheme I, is dissolved in aqueous ethanol and a base such as 10% NaOH is added. The reaction is heated to reflux to give the acids 36. The acids 36 can be converted to the fused pyrazole with a hydrido radical by decarboxylation by heating to about 290°C to give the decarboxylated products 37. The acids 36 also can be converted to the appropriate amides 38 by dissolving in methanol and treating with an appropriate amine in the presence of a condensing agent such as dicyclohexylcarbodiimide (DCC). The amides 38 can also be prepared by direct aminolysis of 35.
- 10
- 15

SCHEME XII



5 Synthetic Scheme XII shows the two step procedure
for preparation of substituted heteroarylhydrazine
compounds 41 as used in Scheme I where R^2 is thienyl.
In step 1, the heteroarylsulfonyl chloride 39 (where LG
represents a leaving group such as halo) is treated
10 with ammonia to give the heteroaryl sulfonamides 40. In
step 2, the heteroaryl sulfonamides 40 are treated with
hydrazine to give the substituted heteroarylhydrazines
41.

The following examples contain detailed
15 descriptions of the methods of preparation of compounds
of Formula I-II. These detailed descriptions fall
within the scope, and serve to exemplify, the above
described General Synthetic Procedures which form part
of the invention. These detailed descriptions are
20 presented for illustrative purposes only and are not
intended as a restriction on the scope of the
invention. All parts are by weight and temperatures are
in Degrees centigrade unless otherwise indicated.

EXAMPLE 1

5 **4-[1,5-Dihydro-6-fluoro-7-methoxy-3-**
 (trifluoromethyl)-[2]benzothiopyrano[4,3-
 c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 2-fluoro-3-methoxybenzoic acid.

- 10 A mixture of potassium tert-butoxide (30.80 g, 274 mmol) and anhydrous tetrahydrofuran (300 mL) was cooled to -78°C and treated with a 1.6 M solution of n-butyllithium (172 mL, 275 mmol) in hexanes. After stirring for 15 minutes, 2-fluoroanisole (31.35 g, 248 mmol) was added and the reaction was stirred an additional 1.8 hours. The reaction was poured into dry ice and warmed to room temperature. Water (250 mL) was added and after extracting with ether (160 mL), the aqueous layer was acidified with concentrated
- 15 hydrochloric acid, and filtered to give 2-fluoro-3-methoxybenzoic acid (21.43 g, 51%) as a yellow solid: mp 155-160°C; ¹H NMR (acetone-*d*₆) 300 MHz 7.46 (ddd, J=6.0 Hz J=1.8 Hz J=1.4 Hz, 1H) 7.36 (dt, J=1.6 Hz J=8.1 Hz, 1H) 7.20 (dt, J=1.4 Hz J=8.1 Hz, 1H) 3.92
- 20 (s, 3H); ¹⁹F NMR (acetone-*d*₆) 300 MHz -134.04 (m). Mass spectrum: M+H=171.

Step 2. Preparation of 2-fluoro-3-methoxybenzyl alcohol.

- 30 2-Fluoro-3-methoxybenzoic acid from Step 1 (16.65 g, 98 mmol) was dissolved in anhydrous

- tetrahydrofuran (60 mL), cooled in an ice bath, and treated with borane dimethyl sulfide complex (19 mL, 190 mmol). The reaction was stirred at room temperature for 4.2 hours, quenched by the slow addition of methanol, and concentrated in vacuo. The residue was dissolved in ethyl acetate, treated with 3N hydrochloric acid and filtered through diatomaceous earth. The organic layer of the filtrate was collected, washed with NaHCO₃, brine, dried over MgSO₄ and reconstituted in vacuo to give 2-fluoro-3-methoxybenzyl alcohol (12.35 g, 81%) as a white solid: mp 53-57°C; ¹H NMR (acetone-d₆) 300 MHz 7.07 (m, 3H) 4.67 (d, J=5.8 Hz, 2H) 4.24 (t, J=5.8 Hz, 1H) 3.86 (s, 3H); ¹⁹F NMR (acetone-d₆) 300 MHz -144.77(m).

15

Step 3. Preparation of 2-fluoro-3-methoxybenzyl chloride.

- 2-Fluoro-3-methoxybenzyl alcohol from Step 2 (12.16 g, 78 mmol) was dissolved in concentrated hydrochloric acid (60 mL) and hydrochloric acid gas was bubbled through the solution for 3 minutes. The reaction was stirred at room temperature (21 hours). The reaction mixture was extracted with ether, dried over MgSO₄ and concentrated in vacuo to give 2-fluoro-3-methoxybenzyl chloride (10.36 g, 76%) as a green oil: ¹H NMR (acetone-d₆) 300 MHz 7.12 (m, 2H) 7.05 (m, 1H) 4.73 (s, 2H) 3.89 (s, 3H); ¹⁹F NMR (acetone-d₆) 300 MHz -142.07(m).

30 Step 4. Preparation of S-(2-fluoro-3-methoxybenzyl)-isothiuronium chloride.

- Thiourea (4.47 g, 59 mmol) was added to a solution of 2-fluoro-3-methoxybenzyl chloride from Step 3 (10.20 g, 58 mmol) in methanol (25 mL). The reaction was heated to reflux for 3.3 hours, concentrated in vacuo, triturated with ether, and filtered to give a white solid (14.65 g, 100%).

Step 5. Preparation of 3-(2-fluoro-3-methoxyphenylthio)propanoic acid.

The thiouronium salt from Step 4 (14.65 g, 58 mmol) was added to sodium chloroacetate (10.44 g, 90 mmol), ethanol (60 mL) and water (25 mL). After heating to reflux, a solution of NaOH (10.66 g, 266 mmol) in water (35 mL) was added to the reaction dropwise. After stirring for 16.6 hours, the reaction was acidified with concentrated hydrochloric acid, extracted with ether, washed with brine, dried over MgSO₄, concentrated in vacuo and recrystallized from ether/hexane to give a brown solid (7.70 g, 57%): mp 72-74°C; ¹H NMR (CDCl₃) 300 MHz 7.03 (m, 1H) 6.91 (m, 2H) 3.89 (d, J=1.0 Hz, 2H) 3.88 (s, 3H) 3.19 (s, 2H); ¹⁹F NMR (CDCl₃) 300 MHz -140.76(m).

Step 6. Preparation of 8-fluoro-7-methoxyisothiochroman-4-one.

The acid from Step 5 (7.63 g, 33 mmol) was dissolved in trifluoroacetic acid (12 mL), treated with trifluoroacetic anhydride (4 mL) and stirred at room temperature (8 minutes). The reaction was poured into 10% Na₂CO₃ (50 mL) and extracted with ethyl acetate, washed with brine, dried over MgSO₄ and concentrated in vacuo to give 8-fluoro-7-methoxyisothiochroman-4-one (5.42 g, 77%) as a brown solid: mp 85-92°C; ¹H NMR (CDCl₃) 300 MHz 7.83 (d, J=8.9 Hz, 1H) 7.19 (t, J=8.5 Hz, 1H) 4.01 (s, 2H) 3.98 (s, 3H) 3.55 (s, 2H); ¹⁹F NMR (CDCl₃) 300 MHz -141.24(m).

Step 7. Preparation of 6-fluoro-7-methoxy-3-(trifluoroacetyl)isothiochroman-4-one.

8-Fluoro-7-methoxyisothiochroman-4-one from Step 6 (1.70 g, 8.0 mmol) was dissolved in anhydrous tetrahydrofuran (30 mL), cooled to -78°C, and treated with a 1.0M tetrahydrofuran solution of sodium bistrimethylsilyl amide (10 mL, 10mmol). After 30


minutes, N-trifluoroacetylimidazole (1.85 g in 10.0 mL THF, 11.3 mmol) was added and the reaction was stirred and warmed to room temperature overnight (19.4 hours). The reaction was treated with 1N hydrochloric acid (30 mL). The organic layer was collected, washed with brine, dried over MgSO_4 , concentrated in vacuo and recrystallized from dichloromethane/isooctane to give the diketone (1.14 g, 46%) as a brown solid: mp 162-164°C; ^1H NMR (CDCl_3) 300 MHz 15.50 (s, 1H) 7.82 (dd, J=8.9 Hz J=1.0 Hz, 1H) 6.99 (t, J=8.5 Hz, 1H) 3.99 (s, 3H) 3.89 (s, 2H) 2.43 (s, 2H); ^{19}F NMR (CDCl_3) 300 MHz: -72.46 (s) -140.30 (d). Mass spectrum: M+H=309.

15 Step 8. Preparation of 4-[6-fluoro-1,5-dihydro-7-methoxy-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (0.77 g, 3.4 mmol) was added to a stirred solution of the diketone from Step 7 (0.93 g, 3.0 mmol) in ethanol (10 mL). The reaction was heated to reflux and stirred for 20.4 hours. The reaction mixture was filtered while hot to give the pyrazole as gray needles (0.55 g, 40%): mp 250-252°C; ^1H NMR (acetone- d_6) 300 MHz 8.08 (d, J=8.9 Hz, 2H) 7.85 (d, J=8.7 Hz, 2H) 7.01 (t, J=8.7 Hz, 1H) 6.81 (br s, 2H) 6.72 (dd, J=8.6 Hz J=1.9 Hz, 1H) 4.20 (s, 2H) 3.91 (s, 3H); ^{19}F NMR (acetone- d_6) 300 MHz -63.32 (s) -140.16 (d).

30

EXAMPLE 2



1,5-Dihydro-6-fluoro-7-methoxy-1-[(4-methylsulfonyl)phenyl]-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazole

5

4-(Methylsulfonyl)phenylhydrazine (0.75 g, 4.0 mmol) was converted to the hydrochloride salt by stirring with a 4N solution of hydrochloric acid in dioxane (10 mL) for 30 minutes. The dioxane was removed in vacuo and the 4-(methylsulfonyl)phenyl hydrazine hydrochloride was combined with the diketone from Example 1, Step 7 (0.89 g, 2.9 mmol) and ethanol (15 mL), heated to reflux and stirred for 14.5 hours. The reaction mixture was filtered while hot and the filtrate was concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with water and with brine, dried over MgSO₄, reconcentrated in vacuo and passed through a column of silica gel, eluting with 20% ethyl acetate/hexane to give the pyrazole (0.46 g, 35%) as a yellow solid: mp 213-215°C; ¹H NMR (acetone-d₆) 300 MHz 8.15 (d, J=8.7 Hz, 2H) 7.94 (d, J=8.7 Hz, 2H) 7.01 (t, J=8.7 Hz, 1H) 6.73 (d, J=8.7 Hz, 1H) 4.21 (s, 2H) 3.90 (s, 3H) 3.23 (s, 3H); ¹⁹F NMR (acetone-d₆) 300 MHz -63.41 (s) -140.17 (d). Mass spectrum: M+=458.1.


10

15

20

25

EXAMPLE 3



30


**4-[1,4-Dihydro-3-(trifluoromethyl)-[1]
benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide**

5 Step 1. Preparation of 3-(trifluoroacetyl)-4-chromanone.

Ethyl trifluoroacetate (9.78 g, 68 mmol) was dissolved in ether (50 mL). To the stirred solution was added 25% sodium methoxide (15.11 g, 70 mmol),
10 followed by 4-chromanone (10.07 g, 68 mmol) dissolved in ether (25 mL). The reaction was stirred at room temperature overnight (18.3 hours), poured into a separatory funnel and washed with 3N hydrochloric acid (20 mL) and with brine (20 mL), dried over MgSO₄,
15 concentrated in vacuo, and recrystallized from ether/hexane to give a yellow solid (10.72 g, 65%): mp 81-83°C; ¹H NMR (CDCl₃) 300 MHz 16.04 (br s, 1H) 7.84 (d, J=7.9 Hz, 1H) 7.51 (m, 1H) 7.09 (m, 1H) 6.98 (d, J=8.5 Hz, 1H) 5.08 (s, 2H); ¹⁹F NMR (CDCl₃) 300 MHz:
20 -72.56 (s). Mass spectrum: M⁺=244.

Step 2. Preparation of 4-[1,4-dihydro-3-(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

25 4-Sulfonamidophenylhydrazine hydrochloride (4.57 g, 20.4 mmol) was added to a stirred solution of the diketone from Step 1 (4.59 g, 18.8 mmol) in ethanol (80 mL). The reaction was heated to reflux and stirred overnight (17.3 hours). The reaction mixture
30 was filtered while hot to give the pyrazole as a white solid (3.99 g, 54%). Upon cooling, the filtrate yielded an additional 1.97 g (26%): mp 250-251°C; ¹H NMR (acetone-d₆) 300 MHz 8.14 (d, J=8.7 Hz, 2H), 7.87 (d, J=8.7 Hz, 2H) 7.30 (m, 1H), 7.08 (d, J=8.1 Hz, 1H)
35 6.89 (m, 3H) 5.41 (s, 2H); ¹⁹F NMR (acetone-d₆) 300 MHz -62.42 (s). High resolution mass spectrum Calc'd. for C₁₇H₁₂F₃N₃O₃S: 395.0551. Found: 395.0551.

EXAMPLE 4

5 **Methyl [1-[4-(aminosulfonyl)phenyl]-1,4-dihydro-[1]benzopyrano[4,3-c]pyrazol-3-yl]carboxylate**

10 Step 1. Preparation of methyl-3-(1-oxo-2-carboxyl)-4-chromanone.

4-Chromanone (9.72 g, 65.6 mmol) was added to dimethyl oxalate (8.92 g, 75.5 mmol) and methanol (75 mL). The solution was treated with sodium methoxide (25%) in methanol (18.52 g, 85.7 mmol) and stirred at room temperature for 18.8 hours. The reaction was treated with 3N hydrochloric acid (30 mL), filtered, and recrystallized from ethyl acetate/isooctane to give the diketone (11.31 g, 74%) as a yellow solid: mp 85-87°C; ¹H NMR (acetone-d₆) 300 MHz 7.87 (d, J=7.9 Hz, 1H) 7.61 (m, 1H) 7.14 (m, 1H) 7.02 (d, J=8.3 Hz, 1H) 5.35 (s, 2H) 3.92 (s, 3H). Mass spectrum: M+Li=234.


25 Step 2. Preparation of methyl [1-[4-(aminosulfonyl)phenyl]-1,4-dihydro-[1]benzopyrano[4,3-c]pyrazol-3-yl]carboxylate.

4-Sulfonamidophenylhydrazine hydrochloride (6.52 g, 29.1 mmol) was added to a stirred solution of the diketone from Step 1 (6.23 g, 26.6 mmol) in methanol (MeOH) (150 mL). The reaction was heated to reflux and stirred for 15.1 hours. The reaction mixture was filtered, washed with MeOH and dried under vacuum to

100

give the pyrazole as a pale green solid (9.81 g, 96%):
mp > 304°C; ¹H NMR (DMSO-*d*₆) 300 MHz 8.02 (d, J=8.7
Hz, 2H) 7.83 (d, J=8.5 Hz, 2H) 7.60 (br s, 2H) 7.25
(2d, 1H) 7.06 (d, J=7.5 Hz, 1H) 6.84 (2d, 1H) 6.71 (d,
5 J=7.9 Hz, 1H) 5.46 (s, 2H) 3.85 (s, 3H). Mass
spectrum: M+H = 386.

EXAMPLE 5



10
4-[1,4-Dihydro-8-fluoro-3-(trifluoromethyl)-
[1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide


15 Step 1. Preparation of 6-fluoro-3- (trifluoroacetyl)-4-chromanone.

Ethyl trifluoroacetate (4.33 g, 30 mmol) was
dissolved in ether (25 mL) and treated with sodium
20 methoxide (25%, 7.08 g, 33 mmol). To the stirred
solution was added 6-fluoro-4-chromanone (4.92 g, 30
mmol) and additional ether (10mL). The reaction was
stirred at room temperature overnight (19.0 hours) and
treated with 3N hydrochloric acid (15 mL). The
25 organic layer was collected, washed with brine, dried
over MgSO₄, concentrated in vacuo and recrystallized
from ether/hexane to give a yellow solid (3.98 g,
51%): mp 107-112°C; ¹H NMR (CDCl₃) 300 MHz 14.95 (s,
1H) 7.52 (dd, 1H) 7.23 (m, 1H) 6.97 (m, 1H) 5.07 (s,
30 2H); ¹⁹F NMR (CDCl₃) 300 MHz -72.60 (s), -119.93 (m).
Mass spectrum: M+=262.

Step 2. Preparation of 4-[1,4-dihydro-8-fluoro-3-(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (1.54 g, 6.9 mmol) was added to a stirred solution of the diketone from Step 1 (1.62 g, 6.2 mmol) in ethanol (35 mL). The reaction was heated to reflux and stirred overnight (17.3 hours). The reaction mixture was filtered while hot to give the pyrazole as a white solid (1.09 g, 43%). Upon cooling, the filtrate yielded an additional 0.70 g (27%): mp 251-251.5°C; ¹H NMR (acetone-*d*₆) 300 MHz 8.17 (d, J=8.5 Hz, 2H) 7.91 (d, J=8.7 Hz, 2H) 7.11 (m, 2H) 6.87 (br s, 1H) 6.58 (dd, J=2.4Hz, 9.5Hz, 3H) 5.41 (s, 2H); ¹⁹F NMR (acetone-*d*₆) 300 MHz -62.46 (s). High resolution mass spectrum Calc'd. for C₁₇H₁₁F₄N₃O₃S: 413.0457. Found: 413.0462.

EXAMPLE 6



4-[3-(Trifluoromethyl)-1H-benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 2-(trifluoroacetyl)-3-coumaranone.

Ethyl trifluoroacetate (1.90 g, 14 mmol) was dissolved in ether (15 mL) and treated with sodium methoxide (25%) (3.67 g, 17 mmol). To the stirred solution was added 3-coumaranone (1.50 g, 11 mmol). The reaction was stirred at room temperature overnight


(19 hours) and treated with 3N HCl (8 mL). The organic layer was collected, washed with brine, dried over MgSO₄, and concentrated *in vacuo* to give a reddish brown solid (2.19 g, 85%): mp 108-111°C; ¹H NMR (CDCl₃) 300 MHz 7.84 (d, J=8.1 Hz, 1H) 7.66 (m, 1H) 7.52 (d, J=8.7 Hz, 1H) 7.37 (m, 1H). Mass spectrum: M+H=231.

10 Step 2. Preparation of 4-[3-(trifluoromethyl)-1H-benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (1.15 g, 5.0 mmol) was added to a stirred solution of the diketone from Step 1 (1.13 g, 4.9 mmol) in ethanol (10 mL). The reaction was heated to reflux and stirred for 2.5 hours. The reaction mixture was filtered to give the pyrazole as a brown solid (1.07 g, 57%): mp 190-195°C; ¹H NMR (acetone-*d*₆) 300 MHz 7.82-7.90 (m, 3H) 7.34-7.53 (m, 5H); ¹⁹F NMR (acetone-*d*₆) 300 MHz -65.47 (s).

20

EXAMPLE 7



25 4-[1,4-Dihydro-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide

30 Step 1. Preparation of 3-(trifluoroacetyl)thio-4-chromanone.


Ethyl trifluoroacetate (8.78 g, 62 mmol) was dissolved in ether (50 mL). To the stirred solution

was added sodium methoxide (14.35 g, 66 mmol) followed by thio-4-chromanone (9.43 g, 57 mmol) dissolved in ether (10mL). The reaction was stirred at room temperature overnight (17.8 hrs) and treated with 3N HCl (25 mL). The organic layer was collected, washed with brine, dried over MgSO₄, concentrated in vacuo and recrystallized from ether/hexane to give a yellow solid (10.20 g, 68%): mp 75-79°C; ¹H NMR (CDCl₃) 300 MHz 15.62 (s, 1H) 7.99 (d, J=7.9Hz, 1H) 7.24-7.42 (m, 3H) 3.81 (s, 2H); ¹⁹F NMR (CDCl₃) 300 MHz -71.92 (s). Mass spectrum: M⁺ = 260.

Step 2. Preparation of 4-[1,4-dihydro-3-(trifluoromethyl)-1,1-benzothioopyranof[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (5.74 g, 25.6 mmol) was added to a stirred solution of the diketone from Step 1 (6.04 g, 23.2 mmol) in ethanol (95 mL). The reaction was heated to reflux and stirred overnight (17.0 hours). The reaction mixture was filtered, and the filtrate cooled to 0°C and filtered to give the pyrazole as a yellow solid (4.42 g, 46%): mp 213-215°C; ¹H NMR (acetone-d₆) 300 MHz 8.09 (d, J=8.9 Hz, 2H) 7.75 (d, J=8.7 Hz, 2H) 7.53 (d, J=7.8Hz, 1H) 7.29 (2d, 1H) 7.07 (2d, 1H) 6.94 (d, J=8.1Hz, 2H) 6.92 (br s, 1H) 4.09 (s, 2H); ¹⁹F NMR (acetone-d₆) 300 MHz -62.22 (s). High resolution mass spectrum Calc'd. for C₁₇H₁₂F₃N₃O₂S₂: 411.0323. Found: 411.0330.

EXAMPLE 8



5 Methyl 1-[(4-aminosulfonyl)phenyl]-1,4-
 dihydro-[1]benzothiopyrano[4,3-c]pyrazol-3-
 yl carboxylate

10 Step 1. Preparation of methyl-3-(1-oxo-2-
 carboxylthiochroman-4-one.

Thiochroman-4-one (9.84 g, 59.9 mmol) was added to dimethyl oxalate (8.54 g, 72.3 mmol) and methanol (75 mL). The reaction was treated with sodium methoxide (25% in methanol, 15.61 g, 72.2 mmol) and stirred at room temperature for 17.8 hours. The reaction was treated with 3N hydrochloric acid (30 mL) and filtered to give the diketone (13.58 g, 90%) as an orange solid: mp 94-97°C; ¹H NMR (CDCl₃) 300 MHz 15.94 (s, 1H) 8.01 (d, J=7.9 Hz, 1H) 7.24-7.39 (m, 3H) 4.11 (s, 2H) 3.94 (s, 3H). Mass spectrum: M⁺=250.


25 Step 2. Preparation of methyl 1-[4-
 (aminosulfonyl)phenyl]-1,4-dihydro-
 [1]benzothiopyrano[4,3-c]pyrazol-3-yl carboxylate.

4-Sulfonamidophenylhydrazine hydrochloride (12.06 g, 53.9 mmol) was added to a stirred solution of the diketone from Step 1 (12.23 g, 48.9 mmol) in MeOH (250 mL). The reaction was heated to reflux and stirred for 6.5 hours. The reaction mixture was filtered, washed with MeOH and dried under vacuum to give the pyrazole as an orange solid (17.88 g, 91%): mp 265-269°C; ¹H NMR (DMSO-*d*₆) 300 MHz 7.97 (d, J=8.5

105

Hz, 2H) 7.69 (d, J=8.5 Hz, 2H) 7.58 (br s, 2H) 7.50 (d, J=7.9 Hz, 1H) 7.24 (t, J=7.7 Hz, 1H) 7.04 (d, J=7.7 Hz, 1H) 6.76 (d, J=7.9 Hz, 1H) 4.21 (s, 2H) 3.86 (s, 3H). Mass spectrum: M+H=402.

5

EXAMPLE 9

10

4-[6,7-Dichloro-1,4-dihydro-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 3-(2,3-15 dichlorophenylthio)propanoic acid.

2,3-Dichlorothiophenol (4.94 g, 28 mmol) was added to acrylic acid (2.11 g, 29 mmol) and stirred at 50°C for 3 hours. The reaction mixture was poured into 10% Na₂CO₃ and extracted with ether. The aqueous layer was acidified with concentrated hydrochloric acid, extracted with ether, washed with brine, dried over MgSO₄, and concentrated in vacuo to give the 3-(2,3-dichlorophenylthio)propanoic acid (3.88 g), contaminated with some acrylic acid, as a clear oil which was used without further purification in the next step.

Step 2. Preparation of 7,8-dichlorothiochroman-4-one.

30 The 3-(2,3-dichlorophenylthio)propanoic acid from Step 1 (3.88 g, 15 mmol) was dissolved in

trifluoroacetic acid (10 mL), treated with trifluoroacetic anhydride (5 mL) and stirred at room temperature for 68.2 hours. The reaction mixture was poured into 10% Na₂CO₃ (100 mL), extracted with ethyl acetate, washed with brine, dried over MgSO₄, and concentrated *in vacuo* to give a yellow oil. The crude material was passed through a column of silica gel eluting with 40% ethyl acetate/hexane to give 7,8-dichlorothiochroman-4-one as a white solid (0.46 g, 13%): mp 93-102°C; ¹H NMR (CDCl₃) 300 MHz 7.99 (d, J=8.7 Hz, 1H) 7.27 (d, J=8.7 Hz, 1H) 3.25 (m, 2H) 2.97 (m, 2H). Mass spectrum: M⁺=233.

Step 3. Preparation of 7,8-dichloro-3-(trifluoroacetyl)thiochroman-4-one.

Ethyl trifluoroacetate (0.29 g, 2.0 mmol) was dissolved in ether (12 mL). To the stirred solution was added sodium methoxide (25%) (0.84 g, 3.9 mmol), followed by 7,8-dichlorothiochroman-4-one from Step 2 (0.42 g, 1.8 mmol). The reaction was stirred at room temperature overnight (19.4 hours) and treated with 3N hydrochloric acid. The organic layer was collected, washed with brine, dried over MgSO₄, and concentrated *in vacuo* to give a brown oily solid which was used without purification in the next step.


Step 4. Preparation of 4-[6,7-dichloro-1,4-dihydro-3-(trifluoromethyl)-1H-benzothiopyran[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (0.32 g, 1.4 mmol) was added to a stirred solution of the diketone from Step 3 (0.44 g, 1.3 mmol) in ethanol (10 mL). The reaction was heated to reflux and stirred overnight (19.4 hours). The reaction mixture was filtered and the filtrate concentrated *in vacuo*, dissolved in ethyl acetate, washed with water and brine, dried over MgSO₄, reconcentrated *in vacuo*, and passed through a column of silica gel (hexane/ethyl

acetate) to give the pyrazole as a white solid (0.24 g, 38%): ^1H NMR (acetone- d_6) 300 MHz 8.08 (d, $J=8.7$ Hz, 2H) 7.77 (d, $J=8.7$ Hz, 2H) 7.27 (d, $J=8.5$ Hz, 1H) 6.95 (d, $J=8.5$ Hz, 1H) 6.79 (br s, 2H) 4.23 (s, 2H);

5 ^{19}F NMR (acetone- d_6) 300 MHz -62.25 (s). High resolution mass spectrum Calc'd. for $\text{C}_{17}\text{H}_{10}\text{Cl}_2\text{F}_3\text{N}_3\text{O}_2\text{S}_2$: 479.9622. Found: 479.9565.

EXAMPLE 10



4-[1,4-Dihydro-7-fluoro-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-
15 yl]benzenesulfonamide

Step 1. Preparation of 3-(3-fluorophenylthio)propanoic acid.

3-Fluorothiophenol (5.39 g, 42 mmol) was added
20 to acrylic acid (3.73 g, 52 mmol) and stirred at room temperature for 20.2 hours. The reaction mixture solidified, was dissolved in ether and extracted with 10% Na_2CO_3 . The aqueous layer was acidified with concentrated hydrochloric acid, extracted with ether,
25 washed with brine, dried over MgSO_4 , and concentrated in vacuo to give the 3-(3-fluorophenylthio)propanoic acid (6.75 g), contaminated with some acrylic acid, as a white solid which was used without further purification in the next step.

30

Step 2. Preparation of 7-fluorothiochroman-4-one.

The acid from Step 1 (6.75 g, 34 mmol) was dissolved in trifluoroacetic acid (20 mL), treated with trifluoroacetic anhydride (10 mL) and stirred at room temperature for 2.2 hours. The reaction mixture was poured into 10% Na₂CO₃ (100 mL), extracted with ether, washed with brine, dried over MgSO₄, and concentrated in vacuo to give a yellow oil which was passed through a column of silica gel with 20% ethyl acetate/hexane to give 7-fluorothiochroman-4-one as a white solid (2.09 g, 34%): mp 61-66°C; ¹H NMR (CDCl₃) 300 MHz 8.13 (m, 1H) 6.98 (m, 1H) 6.86 (m, 1H) 3.23 (m, 2H) 2.99 (m, 2H); ¹⁹F NMR (CDCl₃) 300 MHz -104.70 (m). Mass spectrum: M+H=183.

15 Step 3. Preparation of 7-fluoro-3-(trifluoroacetyl)thiochroman-4-one.

Ethyl trifluoroacetate (1.04 g, 7.3 mmol) was added to solution of 7-fluorothiochroman-4-one from Step 2 (1.24 g, 6.8 mmol) in ether (15 mL). The reaction was treated with 25% sodium methoxide (1.86 g, 8.6 mmol) and stirred at room temperature for 20.9 hours, then treated with 3N hydrochloric acid (10 mL). The organic layer was collected, washed with brine, dried over MgSO₄, concentrated in vacuo, and recrystallized from dichloromethane/isooctane to give the diketone as a yellow solid (0.58g, 31%): mp 84-89°C; ¹H NMR (CDCl₃) 300 MHz 15.65 (s, 1H) 8.03 (m, 1H) 7.08 (m, 1H) 6.97 (m, 1H) 3.83 (s, 2H); ¹⁹F NMR (CDCl₃) 300 MHz -71.81 (s) -103.04 (m).

30


Step 4. Preparation of 4-[1,4-dihydro-7-fluoro-3-(trifluoromethyl)-(1-benzothiopyrano[4,3-c]pyrazol-1-yl)benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (1.36 g, 6.1 mmol) was added to a stirred solution of the diketone from Step 3 (1.65 g, 5.9 mmol) in ethanol (10 mL). The reaction was heated to reflux and stirred overnight (15.2 hours). The reaction mixture was

109

filtered and the filtrate cooled in ice to give the pyrazole as a white solid (0.24 g, 38%): ¹H NMR (acetone-d₆) 300 MHz 8.09 (d, J=8.7 Hz, 2H) 7.76 (d, J=8.7 Hz, 2H) 7.38 (d, J=9.1Hz, 1H) 6.99 (m, 1H) 6.88 (m, 1H) 6.80 (br s, 2H) 4.14 (s, 2H); ¹⁹F NMR (acetone-d₆) 300 MHz: -62.25 (s) -112.68 (m). High resolution mass spectrum Calc'd. for C₁₇H₁₁F₄N₃O₂S₂: 429.0229. Found: 429.0205.

10

EXAMPLE 11

4-[1,4-Dihydro-6-isopropyl-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 3-(2-isopropylphenylthio)propanoic acid.

20 2-Isopropylthiophenol (4.77 g, 31 mmol) was placed in a flask with acrylic acid (2.37 g, 33 mmol) and stirred at room temperature for 71.8 hours. The reaction mixture solidified, was dissolved in ethyl acetate and extracted with 5% NaOH. The aqueous
25 layer was acidified with concentrated hydrochloric acid, extracted with ethyl acetate, dried over MgSO₄, and concentrated in vacuo to give 3-(2-isopropylphenylthio)propanoic acid (6.85 g), contaminated with some acrylic acid, as a yellow

solid which was used without further purification in the next step.

Step 2. Preparation of 8-isopropylthiochroman-4-one.

- 5 3-(2-Isopropylphenylthio)propanoic acid from
Step 1 (6.85 g, 30 mmol) was dissolved in
trifluoroacetic acid (20 mL), treated with
trifluoroacetic anhydride (12mL) and stirred at room
temperature (64.2 hours). The reaction was
10 concentrated in vacuo, and the residue dissolved in
ethyl acetate, extracted with 5% NaOH, washed with
brine, dried over MgSO₄, and concentrated in vacuo to
give a brown oil which was passed through a column of
silica gel eluted with 12% ether/hexane to give 8-
15 isopropylthiochroman-4-one as a brown oil (1.05 g,
17%): ¹H NMR (CDCl₃) 300 MHz 8.01 (d, J=7.0 Hz, 1H)
7.38 (d, J=7.5 Hz, 1H) 7.17 (t, J=7.8 Hz, 1H) 3.19
(m, 3H) 2.96 (m, 2H) 1.25 (d, J=7.0 Hz 6H).

20 Step 3. Preparation of 8-isopropyl-3-
(trifluoroacetyl)thiochroman-4-one.

- Ethyl trifluoroacetate (0.69 g, 4.9 mmol) was
added to a solution of 2-isopropylthiochroman-4-one
from Step 2 (0.97 g, 4.7 mmol) in ether (10 mL). The
25 reaction was treated with 25% sodium methoxide (1.08
g, 5.0 mmol), stirred at room temperature for 18.2
hours and treated with 3N hydrochloric acid (5 mL).
The organic layer was collected, washed with brine,
dried over MgSO₄ and concentrated in vacuo to give
30 the diketone as a brown oil (0.80 g) which was used
without further purification in the next step.

Step 4. Preparation of 4-[1,4-dihydro-6-isopropyl-3-
(trifluoromethyl)-[1]benzothioipyran[4,3-c]pyrazol-1-
35 yl]benzenesulfonamide.


4-Sulfonamidophenylhydrazine hydrochloride (0.69
g, 3.1 mmol) was added to a stirred solution of the

111

diketone from Step 3 (0.83 g, 2.7 mmol) in ethanol (10 mL). The reaction was heated to reflux, stirred overnight (16.1 hours) and concentrated in vacuo. The residue was dissolved in ethyl acetate washed with water, washed with brine, dried over MgSO₄, re-concentrated in vacuo and passed through a column of silica gel with 20% ethyl acetate/hexane to give the pyrazole as a brown solid (0.43 g, 35%): mp 183-186°C; ¹H NMR (acetone-d₆) 300 MHz 8.10 (d, J=8.7 Hz, 2H) 7.71 (d, J=8.5 Hz, 2H) 7.35 (d, J=7.9 Hz, 1H) 7.04 (t, J=7.9 Hz, 1H) 6.79 (m, 3H) 4.04 (s, 2H) 3.46 (m, 1H) 1.28 (d, J=6.8 Hz, 6H); ¹⁹F NMR (acetone-d₆) 300 MHz -62.19 (s) -112.68 (m). High resolution mass spectrum Calc'd. for C₂₀H₁₈F₃N₃O₂S₂: 453.0793.

Found: 453.0848.

EXAMPLE 12



4-[1,4-Dihydro-7,8-dimethoxy-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 3-(3,4-dimethoxyphenylthio)propanoic acid.

3,4-Dimethoxythiophenol (5.00 g, 29 mmol) was placed in a flask with acrylic acid (2.27 g, 32 mmol) and stirred at room temperature for 22.4 hours. The reaction mixture solidified, was dissolved in ethyl acetate and extracted with 10% Na₂CO₃. The aqueous

- layer was acidified with concentrated hydrochloric acid, extracted with ether, dried over MgSO_4 , and concentrated in vacuo to give 3-(3,4-dimethoxyphenylthio)propanoic acid (5.39 g, 76%),
- 5 contaminated with some acrylic acid, as a white solid which was used without further purification in the next step: mp 62-64°C; ^1H NMR (CDCl_3) 300 MHz 9.80 (br s, 1H) 7.01 (d, $J=8.3$ Hz, 1H) 6.98 (s, 1H) 6.82 (d, $J=8.3$ Hz, 1H) 3.87 (s, 3H) 3.86 (s, 3H) 3.06 (t,
- 10 $J=7.3$ Hz, 2H) 2.63 (t, $J=7.3$ Hz, 2H). Mass spectrum: $M^+=242$.

Step 2. Preparation of 6,7-dimethoxythiochroman-4-one.

- 15 The acid from Step 1 (5.23 g, 22 mmol) was dissolved in trifluoroacetic acid (20 mL), treated with trifluoroacetic anhydride (12 mL) and stirred at room temperature (10 minutes). The reaction was poured into 10% Na_2CO_3 (100 mL) and filtered to
- 20 collect 6,7-dimethoxythiochromanone as a yellow solid (1.12 g, 23%). The filtrate was washed with brine, dried over MgSO_4 , concentrated in vacuo and recrystallized from ethyl acetate/hexane to give more 6,7-dimethoxythiochromanone (1.77 g, 37%) as an
- 25 orange solid: mp 138-143°C; ^1H NMR (CDCl_3) 300 MHz 7.60 (s, 1H) 6.68 (s, 1H) 3.91 (s, 3H) 3.89 (s, 3H) 3.19 (m, 2H) 2.93 (m, 3H). Mass spectrum: $M^+=224$.

Step 3. Preparation of 6,7-dimethoxy-3-

- 30 trifluoroacetylthiochroman-4-one.

- Ethyl trifluoroacetate (0.84 g, 5.9 mmol) was dissolved in tetrahydrofuran (20 mL), and treated with 25% sodium methoxide (1.68 g, 7.8 mmol). To the stirred solution was added 6,7-dimethoxythiochroman-
- 35 4-one from Step 2 (1.12 g, 5.0 mmol). The reaction was stirred at room temperature for 87.5 hours, treated with 3N HCl (25 mL) and filtered to give an

113


orange solid (1.04 g, 65%): mp 148-151°C; ^1H NMR (CDCl₃) 300 MHz 16.05 (s, 1H), 7.48 (s, 1H) 6.78 (s, 1H) 3.94 (s, 3H) 3.92 (s, 3H) 3.83 (s, 2H); ^{19}F NMR (CDCl₃) 300 MHz -71.06 (s). Mass spectrum: M+H=

5 321.

Step 4. Preparation of 4-[1,4-dihydro-7,8-dimethoxy-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-clpyrazol-1-yl]benzenesulfonamide.

- 10 4-Sulfonamidophenylhydrazine hydrochloride (0.76 g, 3.4 mmol) was added to a stirred solution of the diketone from Step 3 (1.00 g, 3.1 mmol) in ethanol (20 mL). The reaction was heated to reflux and stirred overnight (15.5 hours). The reaction mixture
- 15 was filtered and the filtrate concentrated *in vacuo*, dissolved in ethyl acetate, washed with water and brine, dried over MgSO₄, reconcentrated *in vacuo*, and recrystallized from ethyl acetate/isooctane to give the pyrazole as a yellow solid (0.72 g, 49%): mp 164-
- 20 168°C; ^1H NMR (acetone-*d*₆) 300 MHz 8.13 (d, J=8.7 Hz, 2H) 7.77 (d, J=8.5 Hz, 2H) 7.06 (s, 1H) 6.82 (br s, 2H) 6.36 (s, 1H) 4.04 (s, 2H) 3.85 (s, 3H) 3.85 (s, 3H); ^{19}F NMR (acetone-*d*₆) 300 MHz -62.20(s). High resolution mass spectrum Calc'd. for
- 25 C₁₉H₁₆F₃N₃O₄S₂: 471.0534. Found: 471.0534.

EXAMPLE 13



**4-[1,4-Dihydro-7-methoxy-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide**

5 Step 1. Preparation of 3-(3-methoxyphenylthio)propanoic acid.

3-Methoxythiophenol (5.70 g, 41 mmol) was placed in a flask with acrylic acid (2.36 g, 33 mmol) and stirred at room temperature for 113.8 hours. The reaction mixture was dissolved in ether and extracted with 10% Na₂CO₃. The aqueous layer was acidified with concentrated HCl, extracted with ether, dried over MgSO₄, and concentrated in vacuo to give the 3-(3-methoxyphenylthio)propanoic acid (2.55 g, 37%) as a white solid: mp 39-42°C; ¹H NMR (CDCl₃) 300 MHz 7.21 (m, 1H) 6.91 (m, 2H) 6.77 (d, J=8.3 Hz, 1H) 3.79 (s, 3H) 3.16 (t, J=7.5 Hz, 2H) 2.68 (t, J=7.3 Hz, 2H).

20 Step 2. Preparation of 7-methoxythiochroman-4-one.

The acid from Step 1 (2.55 g, 12 mmol) was dissolved in trifluoroacetic acid (10 mL), treated with trifluoroacetic anhydride (5 mL) and stirred at room temperature (10 minutes). The reaction was poured into 10% Na₂CO₃ (60 mL), extracted with ether, washed with brine, dried over MgSO₄, and concentrated in vacuo to give a red oil which was passed through a column of silica gel with 50% ether/hexane to give 7-methoxythiochroman-4-one as an orange solid (1.18 g, 51%): mp 49-51°C; ¹H NMR (CDCl₃) 300 MHz 8.06 (d, J=8.9 Hz, 1H) 6.72 (m, 2H) 3.83 (s, 3H) 3.20 (m, 2H) 2.95 (m, 2H). Mass spectrum: M+H=195.

35 Step 3. Preparation of 7-methoxy-3-(trifluoroacetyl)thiochroman-4-one.


7-Methoxythiochroman-4-one from Step 2 (1.13 g, 5.8 mmol) and ethyl trifluoroacetate (0.87 g, 6.1

115

mmol) were dissolved in ether (15 mL), treated with 25% sodium methoxide (2.13 g, 9.9 mmol), stirred at room temperature for 23.0 hours, and treated with 3N HCl. The organic layer was collected, washed with
5 brine, dried over MgSO_4 , concentrated in vacuo and recrystallized from dichloromethane/isooctane to give the diketone as a yellow solid (0.60 g, 35%): mp 93-98°C; ^1H NMR (CDCl_3) 300 MHz 15.92 (s, 1H) 7.96 (d, $J=8.9$ Hz, 1H) 6.82 (m, 2H) 3.87 (s, 3H) 3.82 (s, 2H);
10 ^{19}F NMR (CDCl_3) 300 MHz -71.43 (s). Mass spectrum: $M+H=291$.

Step 4. Preparation of 4-[1,4-dihydro-7-methoxy-3-(trifluoromethyl)-1,1-benzothiopyranol[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (0.46 g, 2.1 mmol) was added to a stirred solution of the diketone from Step 3 (0.57 g, 2.0 mmol) in ethanol (5 mL). The reaction was heated to reflux and stirred
20 overnight (21.5 hrs). The reaction mixture was filtered while hot to give the pyrazole as a yellow solid (0.63 g, 72%): mp 201-206°C; ^1H NMR (acetone- d_6) 300 MHz 8.08 (d, $J=8.7$ Hz, 2H) 7.73 (d, $J=8.7$ Hz, 2H) 7.09 (d, $J=2.6$ Hz, 1H) 6.84 (d, $J=8.7$ Hz, 1H)
25 6.80 (br s, 1H) 6.66 (dd, $J=2.6$ Hz $J=8.9$ Hz, 1H) 4.07 (s, 2H) 3.83 (s, 3H); ^{19}F NMR (acetone- d_6) 300 MHz -62.24 (s). High resolution mass spectrum Calc'd. for $\text{C}_{18}\text{H}_{14}\text{F}_3\text{N}_3\text{O}_3\text{S}_2$: 441.0429. Found: 441.0457.

EXAMPLE 14

**5 4-[1,4-Dihydro-7-methyl-3-(trifluoromethyl)-
 [1]benzothiopyrano[4,3-c]pyrazol-1-
 yl]benzenesulfonamide**

**Step 1. Preparation of 3-(3-
10 methylphenylthio)propanoic acid.**

3-Thiocresol (9.71 g, 78 mmol) was placed in a flask with acrylic acid (5.65 g, 78 mmol) and stirred at room temperature for 62.9 hours. The reaction mixture solidified and was dissolved in ether and extracted with 10% Na₂CO₃. The aqueous layer was acidified with concentrated hydrochloric acid, extracted with ether, washed with brine, dried over MgSO₄ and concentrated in vacuo to give the 3-(3-dimethylphenylthio)propanoic acid (10.13 g, 66%)
15 contaminated with some acrylic acid as a white solid, which was used without further purification in the next step.
20

Step 2. Preparation of 7-methylthiochroman-4-one.

3-(3-Methylphenylthio)propanoic acid from Step 1 (10.12 g, 52 mmol) was dissolved in trifluoroacetic acid (20 mL), treated with trifluoroacetic anhydride (10 mL) and stirred at room temperature for 1.9 hours. The reaction was poured into 10% Na₂CO₃ (100 mL), extracted with ether, washed with brine, dried over MgSO₄, and concentrated in vacuo to give an
25
30

117

orange oil which was passed through a column of silica gel eluting with 8% ether/hexane to give 7-methylthiochroman-4-one (2.97 g, 32%) as a yellow oil: ¹H NMR (CDCl₃) 300 MHz 7.97 (d, J=8.1 Hz, 1H)

5 7.06 (s, 1H) 6.97 (d, J=7.7 Hz, 1H) 3.17 (m, 2H) 2.95 (m, 2H) 2.31 (s, 3H). Mass spectrum: M+H=179.

Step 3. Preparation of 7-methyl-3-(trifluoroacetyl)thiochroman-4-one.


10 7-Methylthiochroman-4-one from Step 2 (2.92 g, 16 mmol) and ethyl trifluoroacetate (2.45 g, 17 mmol) were dissolved in ether (20 mL), treated with 25% sodium methoxide (4.60 g, 21 mmol), stirred at room temperature for 15.0 hours and treated with 3N
15 hydrochloric acid (15 mL). The organic layer was collected, washed with brine, dried over MgSO₄, concentrated in vacuo, and recrystallized from dichloromethane/isooctane to give the diketone as a yellow solid (3.05g, 68%): mp 68-72°C; ¹H NMR
20 (CDCl₃) 300 MHz 15.73 (s, 1H) 7.89 (d, J=8.1 Hz, 1H) 7.17 (s, 1H) 7.09 (d, J=8.1 Hz, 1H) 3.80 (s, 2H) 2.37 (s, 3H); ¹⁹F NMR (CDCl₃) 300 MHz -71.75 (s). Mass spectrum: M+H=275.

25 Step 4. Preparation of 4-[1,4-dihydro-7-methyl-3-(trifluoromethyl)-11benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (1.21 g, 5.4 mmol) was added to a stirred solution of the
30 diketone from Step 3 (1.41 g, 5.1 mmol) in ethanol (20 mL). The reaction was heated to reflux and stirred overnight (15.8 hours). The reaction mixture was filtered and the filtrate concentrated in vacuo, dissolved in ethyl acetate, washed with water and
35 brine, dried over MgSO₄, reconcentrated in vacuo, and recrystallized from ethyl acetate/isooctane to give the pyrazole as a yellow solid (1.14 g, 52%): mp 251-

118

252°C; ^1H NMR (acetone- d_6) 300 MHz 8.08 (d, $J=8.7$ Hz, 2H) 7.73 (d, $J=8.7$ Hz, 2H) 7.36 (s, 1H) 6.79 (m, 4H) 4.06 (s, 2H) 2.29 (s, 3H); ^{19}F NMR (acetone- d_6) 300 MHz -62.22(s). High resolution mass spectrum
 5 Calc'd. for $\text{C}_{18}\text{H}_{14}\text{F}_3\text{N}_3\text{O}_2\text{S}_2$: 425.0480. Found: 425.0470.

EXAMPLE 15

10 4-[7-Chloro-1,4-dihydro-3-(trifluoromethyl)-
 [1]benzothiopyrano[4,3-c]pyrazol-1-
 y]benzenesulfonamide

15 Step 1. Preparation of 3-(3-chlorophenylthio)propanoic acid

Acrylic acid (2.66 g, 37 mmol) and 3-chlorothiophenol (4.85 g, 34 mmol) were dissolved in
 20 ether (15 mL) and stirred at room temperature for 88.0 hours. The reaction mixture solidified, was dissolved in ether and extracted with 5% NaOH. The aqueous layer was acidified with concentrated hydrochloric acid, extracted with ether, washed with
 25 brine, dried over MgSO_4 , and concentrated in vacuo to give the 3-(3-chlorophenylthio)propanoic acid (2.20 g, 34%): ^1H NMR (acetone- d_6) 300 MHz 7.33-7.40 (m, 3H) 7.24 (m, 1H) 3.25 (t, $J=7.1$ Hz, 2H) 2.67 (t, $J=7.1$ Hz, 2H). Mass spectrum: $M+H=217$.

30 Step 2. Preparation of 7-chlorothiochroman-4-one.

3-(3-Chlorophenylthio)propanoic acid from Step 1 (2.18 g, 10 mmol) was dissolved in trifluoroacetic acid (10 mL), treated with trifluoroacetic anhydride (6 mL) and stirred at room temperature for 67.8 hours. The reaction was concentrated in vacuo and the residue dissolved in dichloromethane, extracted with 5% NaOH, dried over MgSO₄, and reconcentrated in vacuo to give a brown oil. The crude oil was passed through a column of silica gel with 10% ether/hexane to give a mixture of 7-chlorothiochroman-4-one and 5-chlorothiochroman-4-one as a yellow oil (1.30 g) which was carried on to the next step without further purification.

15 Step 3. Preparation of 7-chloro-3-(trifluoroacetyl)thiochroman-4-one.

Ethyl trifluoroacetate (1.02 g, 7.2 mmol) was placed in a round bottom flask and dissolved in ether (10 mL). To the stirred solution was added 25% sodium methoxide (1.80 g, 8.3 mmol) followed by 7-chlorothiochroman-4-one from Step 2 (1.30 g, 6.5 mmol). The reaction was stirred at room temperature overnight (17.3 hrs) and treated with 3N hydrochloric acid (5 mL). The organic layer was collected, washed with brine, dried over MgSO₄, concentrated in vacuo, and recrystallized from ether/hexane to give the diketone (0.61 g, 32%) as a yellow solid: mp 64-71°C; ¹H NMR (CDCl₃) 300 MHz 15.53 (s, 1H) 7.92 (d, J=8.5 Hz, 1H) 7.36 (s, 1H) 7.25 (d, J=8.7 Hz, 1H) 3.82 (s, 2H); ¹⁹F NMR (CDCl₃) 300 MHz -71.97 (s). Mass spectrum: M+=294.


35 Step 4. Preparation of 4-[7-chloro-1,4-dihydro-3-(trifluoromethyl)-[1]benzothiopyran[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (0.43 g, 1.9 mmol) was added to a stirred solution of the

120

diketone from Step 3 (0.55 g, 1.9 mmol) in ethanol (6 mL). The reaction was heated to reflux and stirred for 18.9 hours. The reaction mixture was cooled and filtered to give the pyrazole as a yellow solid (0.15 g, 18%): mp 237-238°C; ¹H NMR (acetone-d₆) 300 MHz 8.09 (d, J=8.7 Hz, 2H) 7.77 (d, J=8.7 Hz, 2H) 7.59 (d, J=2.0 Hz, 1H) 7.10 (dd, J=8.5 Hz J=2.0Hz, 1H) 6.96 (d, J=8.5 Hz, 1H) 6.81 (br s, 2H) 4.14 (s, 2H); ¹⁹F NMR (acetone-d₆) 300 MHz -62.25(s). High resolution mass spectrum Calc'd. for C₁₇H₁₁ClF₃N₃O₂S₂: 444.9933. Found: 444.9874.

EXAMPLE 16



4-[1,5-Dihydro-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide

Step 1. Preparation of 3-(trifluoroacetyl)isothiochroman-4-one.

Ethyl trifluoroacetate (3.67 g, 25.8 mmol) was dissolved in tetrahydrofuran (15 mL) and treated with 25% sodium methoxide (6.46 g, 29.9 mmol) followed by a solution of isothiochroman-4-one (4.15 g, 25.3 mmol) in tetrahydrofuran (15 mL). The reaction was stirred at room temperature for 70.8 hours and treated with 3N hydrochloric acid (10 mL). The organic layer was collected, washed with brine, dried over MgSO₄, concentrated in vacuo, and passed through


121

a column of silica gel eluting with 40% ethyl acetate/hexane to give a brown solid (5.40 g, 82%):
¹H NMR (CDCl₃) 300 MHz 15.30 (s, 1H) 7.99 (d, J=7.9 Hz, 1H) 7.54 (m, 1H) 7.43 (m, 1H) 7.23 (m, 1H) 3.81 (s, 3H).

Step 2. Preparation of 4-[[1,5-dihydro-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

- 10 4-Sulfonamidophenylhydrazine hydrochloride (0.81 g, 3.6 mmol) was added to a stirred solution of the diketone from Step 1 (0.83 g, 3.2 mmol) in ethanol (10 mL). The reaction was heated to reflux and stirred 2.1 hours. The reaction mixture was filtered
- 15 and the filtrate concentrated *in vacuo*. The residue was dissolved in ethyl acetate, washed with water and with brine, dried over MgSO₄, reconcentrated *in vacuo*, and passed through a column of silica gel with 40% ethyl acetate to give the pyrazole (0.15 g, 11%):
- 20 mp 230-232°C; ¹H NMR (acetone-*d*₆) 300 MHz 8.09 (d, J=8.7 Hz, 2H) 7.85 (d, J=8.9 Hz, 2H) 7.53 (d, J=7.7 Hz, 1H) 7.40 (t, J=7.5 Hz, 1H) 7.21 (t, J=7.7 Hz, 1H) 6.93 (d, J=7.9 Hz, 1H) 6.79 (br s, 2H) 4.16 (s, 2H); ¹⁹F NMR (acetone-*d*₆) 300 MHz -62.94(s).
- 25 High resolution mass spectrum Calc'd. for C₁₇H₁₂F₃N₃O₂S₂: 411.0323. Found: 411.0324.

EXAMPLE 17



30

4-[1,5-Dihydro-7-methyl-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide

5 Step 1. Preparation of S-(3-methylbenzyl)-
isothiuronium chloride.

Thiourea (26.19 g, 344 mmol) was added to a solution of α -chloro-m-xylene (48.21 g, 343 mmol) in methanol (50 mL). The reaction was heated to reflux and additional methanol (10 mL) was added to dissolve all of the thiourea. After 64.3 hours, the reaction was filtered and dried under vacuum to give a white solid (68.15 g, 92%): mp 182-186°C; ^1H NMR (DMSO- d_6) 300 MHz 9.34 (br s, 4H) 7.22 (m, 3H) 7.12 (m, 1H) 4.48 (s, 2H) 2.27 (s, 3H).

Step 2. Preparation of 3-(3-methylphenylthio)propanoic acid.

A 250 mL flask was charged with the thiuronium salt from Step 1 (10.99 g, 51 mmol), sodium chloroacetate (8.86 g, 76 mmol), ethanol (95 mL) and water (10 mL). After heating to reflux, a solution of NaOH (9.05 g, 226 mmol) in water (50 mL) was added to the reaction dropwise over seven minutes. After stirring for 3.6 hours, the reaction was acidified with concentrated hydrochloric acid, extracted with ether, washed with brine, dried over MgSO_4 and concentrated in vacuo to give a white solid (9.95 g, 100%): mp 73-75.5°C; ^1H NMR (CDCl_3) 300 MHz 7.16 (m, 4H) 3.83 (s, 2H) 3.12 (s, 2H) 2.35 (s, 3H). Mass spectrum: $M^+=196$.

Step 3. Preparation of 7-methylisothiochroman-4-one.

The acid from Step 2 (6.06 g, 31 mmol) was dissolved in trifluoroacetic acid (11 mL), treated with trifluoroacetic anhydride (5 mL) and stirred at room temperature for 0.33 hours. The reaction was

poured into 10% Na₂CO₃ (100 mL), extracted with ether, washed with brine, dried over MgSO₄, concentrated in vacuo, and recrystallized from ether/hexane to give 7-chloroisothiochroman-4-one

- 5 (2.25 g, 41%) as a white solid: mp 79.5-82°C; ¹H NMR (CDCl₃) 300 MHz 7.97 (d, J=8.1 Hz, 1H) 7.17 (d, J=8.1 Hz, 1H) 7.00 (s, 1H) 3.87 (s, 2H) 3.52 (s, 2H) 2.37 (s, 3H). Mass spectrum: M+H=179.

10 Step 4. Preparation of 7-methyl-3-(trifluoroacetyl)isothiochroman-4-one.

Ethyl trifluoroacetate (1.80 g, 12.7 mmol) was placed in a round bottom flask and dissolved in ether (10 mL). To the stirred solution was added 25%


- 15 sodium methoxide (3.90 g, 18.0 mmol) followed by 7-chloroisothiochroman-4-one from Step 3 (2.25 g, 12.6 mmol) dissolved in ether (10 mL) and tetrahydrofuran (10 mL). The reaction was stirred at room temperature for 24.6 hours and treated with 3N
20 hydrochloric acid. The organic layer was collected, washed with brine, dried over MgSO₄, concentrated in vacuo and passed through a column of silica gel with 20% ether/hexane to give the diketone (1.93 g, 56%) as a brown solid: ¹H NMR (CDCl₃) 300 MHz 15.45 (s,
25 1H) 7.88 (d, J=8.1 Hz, 1H) 7.25 (d, J=8.1 Hz, 1H) 7.06 (s, 1H) 3.77 (s, 2H) 2.43 (s, 2H); ¹⁹F NMR (CDCl₃) 300 MHz: -72.76 (s). Mass spectrum: M+H=275.

30 Step 5. Preparation of 4-[1,5-dihydro-7-methyl-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

- 4-Sulfonamidophenylhydrazine hydrochloride (1.68 g, 7.5 mmol) was added to a stirred solution of the
35 diketone from Step 4 (1.93 g, 7.0 mmol) in ethanol (15 mL). The reaction was heated to reflux and stirred for 15.2 hours. The reaction mixture was

124

concentrated in vacuo and the residue dissolved in ethyl acetate, washed with water and with brine, dried over MgSO_4 , reconcentrated in vacuo and recrystallized from ethyl acetate/isooctane to give the pyrazole as a brown solid (1.48 g, 49%): mp 253-255°C; ^1H NMR (acetone- d_6) 300 MHz 8.08 (d, $J=8.7$ Hz, 2H) 7.83 (d, $J=8.7$ Hz, 2H) 7.35 (s, 1H) 7.02 (d, $J=8.1$ Hz, 1H) 6.78 (m, 3H) 4.11 (s, 2H) 2.34 (s, 3H); ^{19}F NMR (acetone- d_6) 300 MHz -62.94(s). High resolution mass spectrum Calc'd. for $\text{C}_{18}\text{H}_{14}\text{F}_3\text{N}_3\text{O}_2\text{S}_2$: 426.0558. Found: 426.05554.

EXAMPLE 18

1,5-Dihydro-1-[4-(methanesulfonyl)phenyl]-7-methyl-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazole


Step 1. Preparation of 1-[4-methanesulfonylphenyl]-7-methyl-3-(trifluoromethyl)-1,5-dihydro-[2]benzothiopyrano[4,3-c]pyrazole.

4-(Methanesulfonyl)phenylhydrazine (1.23 g, 6.6 mmol) was converted to the hydrochloride salt by stirring with a 4N solution of hydrochloric acid in dioxane (10 mL) for 25 minutes. The dioxane was removed in vacuo, and the 4-(methanesulfonyl)phenylhydrazine hydrochloride was combined with the diketone (Example 17, Step 4) (1.12

125

- g, 2.9 mmol) and ethanol (20 mL), heated to reflux and stirred for 15.5 hours. The reaction mixture was filtered while hot and the filtrate was concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with water and with brine, dried over MgSO_4 , reconcentrated in vacuo, and passed through a column of silica gel eluting with 12% ethyl acetate/hexane to give the pyrazole (0.31 g, 18%) as a yellow solid: mp 207-209°C; ^1H NMR (acetone- d_6) 300 MHz 8.14 (d, $J=8.7$ Hz, 2H) 7.92 (d, $J=8.9$ Hz, 2H) 7.35 (s, 1H) 7.35 (d, $J=8.1$ Hz, 1H) 6.83 (d, $J=7.9$ Hz, 1H) 4.11 (s, 2H) 3.23 (s, 3H) 2.34 (s, 3H); ^{19}F NMR (acetone- d_6) 300 MHz -62.97 (s). High resolution mass spectrum Calc'd. for $\text{C}_{19}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{S}_2$: 424.0527.
- Found: 424.0524.

EXAMPLE 19



- 4-[7-Chloro-1,5-dihydro-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

- Step 1. Preparation of S-(3-chlorobenzyl)-isothiuronium chloride.

3-Chlorobenzyl chloride (24.2 g, 0.15 mol) and thiourea (11.4 g, 0.15 mol) were dissolved in methanol (70 mL) and heated to reflux for 16 hours.

The reaction was cooled to room temperature and a precipitate formed. Ether (150 mL) was added to complete the precipitation of compound. The crystals were isolated by filtration and washed with ether (100 mL). After drying in vacuo, 31.9 g (90%) of pure S-(3-chlorobenzyl)-isothiuronium chloride was obtained: ^1H NMR (CD_3OD) δ = 4.43p (s, 2H), 7.36p (s, 3H), 7.47p (s, 1H).

10 Step 2. Preparation of 3-chlorobenzylmercaptoacetic acid

S-(3-Chlorobenzyl)-isothiuronium chloride from Step 1 (11.86 g, 50 mmol) and chloroacetic acid (7.1 g, 75 mmol) were dissolved in ethanol (100 ml) and heated to 80°C in a 4-neck flask equipped with nitrogen inlet, condenser and addition funnel. A solution of NaOH (10 g) in H_2O (100 mL) and ethanol (50 mL) was added dropwise over 1 hour to this hot solution. The reaction was heated to 100°C for 4 hours. The reaction was cooled to room temperature, acidified with concentrated hydrochloric acid (45 mL) and extracted with ether (500 mL). The organic layer was washed with brine (300 ml), dried over MgSO_4 and concentrated in vacuo to yield 10.84 g (100%) of 3-chlorobenzylmercaptoacetic acid which was used without further purification: ^1H NMR (CDCl_3) δ = 3.08p (s, 2H), 3.80p (s, 2H), 7.23p (m, 3H), 7.34p (s, 1H), 9.07p (broad s, 1H).

30 Step 3. Preparation of 7-chloroisothiochroman-4-one.

3-Chlorobenzylmercaptoacetic acid from Step 2 (3.35 g) was dissolved in trifluoroacetic acid (50 mL). Trifluoroacetic acid anhydride (25 mL) was added and the reaction stirred at reflux, under a nitrogen atmosphere for 16 hours. The solution was carefully poured into 10% Na_2CO_3 solution (500 mL) which was stirring vigorously. The organics were

extracted into ether (500 mL) and washed with brine (300 mL), dried over MgSO_4 and concentrated in vacuo. The resulting brown solid was purified by silica gel chromatography eluting with a 0-10% gradient of ethyl acetate in hexane to yield 7-chloroisothiochroman-4-one (1.57 g, 51%): ^1H NMR (CDCl_3) δ = 3.49p (d, 2H, J = 0.8 Hz), 3.8p (s, 2H), 7.16p (m, 1H), 7.3p (dd, 1H, J = 2.0, 8.4 Hz), 7.96p (d, 1H, J = 8.5 Hz).

10 Step 4. Preparation of 7-chloro-3-(trifluoroacetyl)isothiochroman-4-one.


7-Chloroisothiochroman-4-one from Step 3 (0.3 g, 1.5 mmol) was dissolved in tetrahydrofuran (15 mL) and cooled to -78°C . A solution of sodium bistrimethylsilyl amide (1.5 mL of a 1.0M tetrahydrofuran solution) was added and the reaction stirred for 0.5 hours at -78°C . Trifluoroacetyl imidazole (0.21 mL, 1.8 mmol) was added and the reaction was warmed to room temperature and stirred under a nitrogen atmosphere for 16 hours. 1N Hydrochloric acid (100 mL) was added to the reaction followed by extraction with ether (150 mL). The organics were washed with brine (75 mL), dried over MgSO_4 and concentrated in vacuo. The resulting yellow oil was used in the next step without further purification.

30 Step 5. Preparation of 4-[7-chloro-1,5-dihydro-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

A mixture of the diketone from Step 4 (1.51 mmol) and 4-sulphonamidophenylhydrazine hydrochloride (0.41 g, 1.8 mmol) was dissolved in ethanol (50 mL) and heated to reflux for 16 hours. The reaction was concentrated in vacuo and the resulting solid was dissolved in ethyl acetate (200 mL). The organics were washed with water (200 mL) and with brine (150

mL), dried over MgSO_4 and concentrated in vacuo. The resulting oil was chromatographed on silica gel eluting with a gradient of ethyl acetate (from 10 - 50%) in hexane to yield pure tricyclic pyrazole (0.25 g, 40%): mp 241-242°C; ^1H NMR (acetone- d_6) δ = 4.18p (s, 2H), 6.79p (s, 2H), 6.94p (d, 1H, J = 8.5 Hz), 7.26p (dd, 1H, J = 2.2, 8.4 Hz), 7.6p (d, 1H, J = 2.2 Hz), 7.83p (dd, 2H, J = 2.1, 6.9 Hz), 8.1p (dd, 2H, J = 2.1, 6.7 Hz); ^{19}F NMR (acetone- d_6) δ -62.94p (s, 3F).

EXAMPLE 20



15

4-[1,5-Dihydro-7-methoxy-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide

20 Step 1. Preparation of S-(3-methoxyphenylmethyl)-isothiuronium chloride.

A solution of 3-methoxybenzyl chloride (15.65 g, 0.1 mol) and thiourea (7.6 g, 0.1 mol) were dissolved in 40 mL of ethanol and heated to reflux for 16 hours, during this time the isothiuronium salt crystallized. The thiouronium chloride was isolated by filtration and recrystallized from ether and ethanol (21.85 g, 94%, mp 172.5-174.0°C). This material was used directly in the next step.

Step 2. Preparation of 3-methoxybenzylmercaptoacetic acid

The thiouronium chloride from Step 1 (20.00 g, 86 mmol) and chloroacetic acid (11.07 g, 95 mmol) were dissolved in ethanol (100 mL) and heated to 80°C in a 4-neck flask equipped with nitrogen inlet, condenser and addition funnel. A solution of NaOH (10 g) in water (100 mL) and ethanol (50 mL) was added dropwise over 1 hour to the hot solution. The reaction was heated to 100°C for 4 hours. The reaction was cooled to room temperature, acidified with concentrated hydrochloric acid (45 mL), and extracted with ether (500 mL). The organic layer was washed with brine (300 mL), dried over MgSO₄ and concentrated *in vacuo* to yield an oil that was vacuum distilled to provide 16.41 g (90%) of pure acid: bp 160-170°C at 0.2 mm; ¹H NMR (CDCl₃/300 MHz) 3.1p (s, 2H), 3.8p (s, 3H), 3.82p (s, 2H), 6.8p (m, 1H), 6.91p (s, 1H), 6.96p (m, 1H), 7.23p (t, 1H, J = 7.7 Hz), 8.33p (broad s, 1H), 11.1(brs, 1H).

Step 3. Preparation of 7-methoxyisothiochroman-4-one

3-methoxybenzylmercaptoacetic acid from Step 2 (10.82 g) was dissolved in trifluoroacetic acid (50 mL). Trifluoroacetic acid anhydride (25 mL) was added and the reaction stirred under a nitrogen atmosphere for 0.25 hours. At this time, TLC showed no starting material. The solution was carefully poured into 10% Na₂CO₃ solution (500 mL) which was stirring vigorously. The organics were extracted into ether (500 mL) and washed with brine (300 mL), dried over MgSO₄ and concentrated *in vacuo*. The resulting brown solid was purified by silica gel chromatography eluting with 20% ethyl acetate in hexane to yield 7-methoxyisothiochroman-4-one (4.84 g, 49%): ¹H NMR (CDCl₃) δ = 3.48p (t, 2H, J = 0.8

Hz), 3.83p (s, 3H), 3.84p (s, 2H), 6.6p (d, 1H, J = 2.4 Hz), 6.83p (dd, 1H, J = 2.4, 8.9 Hz), 8.0p (d, 1H, J = 8.9 Hz).

5 Step 4. Preparation of 7-methoxy-3-(trifluoroacetyl)-isothiochroman-4-one.

7-Methoxyisothiochroman-4-one from Step 3 (0.58 g, 3.0 mmol) was dissolved in tetrahydrofuran (30 mL) and cooled to -78°C. A solution of sodium
10 bistrimethylsilylamide (3.0 mL of a 1.0 M tetrahydrofuran solution) was added and the reaction stirred for 0.5 hours at -78°C. Trifluoroacetyl imidazole (0.41 mL, 3.6 mmol) was added and the
15 reaction was warmed to room temperature and stirred under a nitrogen atmosphere for 16 hours. At this time, 1N hydrochloric acid (200 mL) was added to the reaction followed by extraction with ether (250 mL). The organics were washed with brine (150 mL), dried over MgSO₄ and concentrated *in vacuo*. The resulting
20 yellow oil (0.42 g, 48%) was used without further purification.


25 Step 5. Preparation of 4-[1,5-dihydro-7-methoxy-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

A solution of the dione from Step 4 (0.42 g, 1.4 mmol) and 4-sulphonamidophenylhydrazine hydrochloride (0.42 g, 1.8 mmol) were dissolved in ethanol (50 mL) and heated to reflux for 16 hours. The reaction was
30 concentrated *in vacuo* and the resulting solid was dissolved in ethyl acetate (200 mL). The organics were washed with H₂O (200 mL) followed by brine (150 mL), dried over MgSO₄ and concentrated *in vacuo*. The resulting oil was chromatographed on silica gel
35 eluting with a gradient of ethyl acetate (from 20-50%) in hexane to yield pure tricyclic pyrazole (0.25 g, 41%): mp 268-270°C; ¹H NMR (acetone-d₆) δ = 3.84p

131

(s, 3H), 4.12p (s, 2H), 6.8p (m, 3H), 7.1p (d, 1H, J = 2.4 Hz), 7.81p (d, 2H, J = 6.6 Hz), 7.82p (s, 2H), 8.08p (dd, 2H, J = 1.9, 6.6 Hz); ^{19}F NMR (acetone- d_6) δ = -62.9p (s, 3F).

5

EXAMPLE 21

10 4-[7-Chloro-1,5-dihydro-3-trifluoromethyl-
thieno[3',2':4,5]thiopyrano[3,2-c]pyrazol-1-
yl]benzenesulfonamide

15 Step 1. Preparation of S-(5-chloro-2-
thienylmethyl)-isothiuronium chloride.

2-Chloro-5-(chloromethyl)thiophene (14.5 g, 87 mmol) and thiourea (6.6 g, 87 mol) were dissolved in methanol (30 mL) and heated to reflux for 16 hours. The reaction was cooled to room temperature and a precipitate formed. Ether (150 mL) was added to complete the precipitation of compound. The crystals were isolated by filtration and washed with ether (100 mL). After drying in vacuo, 19.0 g (90%) of pure S-(5-chloro-2-thienylmethyl)-isothiuronium chloride were obtained: ^1H NMR (CD_3OD) δ = 4.83p (s, 2H), 6.87p (d, 1H, J = 3.8 Hz), 6.96p (d, 1H, J = 3.2 Hz).

Step 2. Preparation of 5-chloro-2-thienylmethylmercaptoacetic acid.

The compound from Step 1 (12.16 g, 50 mmol) and chloroacetic acid (7.1 g, 75 mmol) were dissolved in ethanol (100 mL) and heated to 80°C in a 4-neck flask equipped with nitrogen inlet, condenser and addition funnel. A solution of NaOH (10 g) in water (100 mL) and ethanol (50 mL) was added dropwise over 1 hour to the hot solution. The reaction was heated to 100°C for 4 hours. The reaction was cooled to room temperature, acidified with concentrated hydrochloric acid (45 mL) and extracted with ether (500 mL). The organic layer was washed with brine (300 mL), dried over MgSO₄ and concentrated *in vacuo* to yield 11.14 g (100%) of pure acid which was used without further purification: ¹H NMR (CDCl₃) δ = 3.2p (s, 2H), 3.98p (s, 2H), 6.73p (d, 1H, J = 3.6 Hz), 6.77p (d, 1H, J = 3.8 Hz).

Step 3 Preparation of 6-chloro-5,6-dihydro-4H-thieno[2,3-b]thiopyran-4-one

The acid from Step 2 (4.45 g) was dissolved in trifluoroacetic acid (45 mL). Trifluoroacetic acid anhydride (20 mL) was added and the reaction was stirred under a nitrogen atmosphere for 0.25 hours. At this time, TLC showed no starting material. The solution was carefully poured into 10% Na₂CO₃ solution (600 mL) which was stirring vigorously. The organics were extracted into ethyl acetate (500 mL), washed with brine (300 mL), dried over MgSO₄ and concentrated *in vacuo*. The resulting brown solid was purified by SiO₂ chromatography eluting with 10% ethyl acetate in hexane to yield of pure intermediate (2.5 g, 61%): ¹H NMR (CDCl₃) δ = 3.33p (d, 2H, J = 0.6 Hz), 3.78p (s, 2H), 7.1p (s, 1H).

Step 4 Preparation of 6-chloro-5,6-dihydro-3-trifluoroacetyl-4H-thieno[2,3-b]thiopyran-4-one

The compound from Step 3 (1.03 g, 5.0 mmol) was dissolved in tetrahydrofuran (50 mL) and cooled to -78°C. A solution of sodium bistrimethylsilylamide (5.0 mL of a 1.0 M tetrahydrofuran solution) was added and the reaction stirred for 0.75 hours at -78°C. Trifluoroacetyl imidazole (0.68 mL, 6.0 mmol) was added and the reaction was warmed to room temperature and stirred under a nitrogen atmosphere for 16 hours. At this time, 1N hydrochloric acid (300 mL) was added to the reaction followed by extraction with ether (350 mL). The organics were washed with brine (200 mL), dried over MgSO₄ and concentrated *in vacuo*. The resulting yellow oil was used without further purification.


Step 5 Preparation of 4-[7-chloro-1,5-dihydro-3-trifluoromethyl-thieno[3',2':4,5]thiopyrano[3,2-c]pyrazol-1-yl]benzenesulfonamide

The compound from Step 4 (5.5 mmol) and 4-sulphonamidophenylhydrazine hydrochloride (1.47 g, 6.6 mmol) were dissolved in ethanol (100 mL) and heated to reflux for 16 hours. The reaction was concentrated *in vacuo* and the resulting solid dissolved in ethyl acetate (300 mL). The organics were washed with water (300 mL) followed by brine (200 mL) and were then dried over MgSO₄ and concentrated *in vacuo*. The resulting oil was chromatographed on silica gel eluting with a gradient of ethyl acetate (from 20 - 40%) in hexane. This product was first crystallized from ethyl acetate and isooctane, then from ethanol and water to yield pure 4-[7-chloro-1,5-dihydro-3-trifluoromethyl-thieno[3',2':4,5]thiopyrano[3,2-c]pyrazol-1-yl]benzenesulfonamide [0.35 g, 16% from Step 2]: mp 218-220°C (dec); ¹H NMR (CDCl₃) δ = 4.0p (s, 2H),

134

6.27p (s, 2H), 7.32p (s, 1H), 7.61(d, 2H, J = 7.0 Hz), 8.02p (d, 2H, J = 7.0 Hz); ^{19}F NMR (acetone- d_6) δ = -59.25p (s, 3F).

5

EXAMPLE 22

10 1-[4-(Aminosulfonyl)phenyl]-1,4-dihydro-
 [1]benzothiopyrano[4,3-c]pyrazol-3-
 carbonitrile


15 Step 1. Preparation of 1-[4-(aminosulfonyl)phenyl]-
 1,4-dihydro-[1]benzothiopyrano[4,3-c]pyrazol-3-
 carboxamide.

 The compound from Example 8 (11.31 g, 28.3 mmol)
 was placed in a 500 mL flask with methanol (200 mL),
 anhydrous ammonia was bubbled through the solution,
 the flask was capped and allowed to stand. After 14
20 days the reaction was concentrated *in vacuo* and
 recrystallized from ethyl acetate to give the
 carboxamide as yellow solid (10.14 g, 93%): mp 238-
 242°C; ^1H NMR (acetone- d_6) 300 MHz 8.07 (d, J=8.7
 Hz, 2H) 7.72 (d, J=8.7 Hz, 2H) 7.49 (d, J=7.9 Hz, 1H)
25 7.30 (br s, 1H) 7.23 (dd, 1H) 7.02 (dd, 1H) 6.91 (d,
 J=7.9 Hz, 1H) 6.80 (br s, 2H) 6.75 (br s, 1H) 4.28
 (s, 2H). Mass spectrum: M+H=387.

30 Step 2. Preparation of 1-[4-(aminosulfonyl)phenyl]-
 1,4-dihydro-[1]benzothiopyrano[4,3-c]pyrazol-3-
 carbonitrile

Dimethylformamide (10 mL) (DMF) was placed in a 250 mL flask and cooled to 0°C. Oxalyl chloride (2 mL, 23 mmol) was added and stirred for 15 minutes. A solution of the product from Step 1 (3.34 g, 9 mmol) in DMF (13 mL) was added and the reaction was stirred for 0.8 hours, treated with pyridine (3.8 mL, 47 mmol), poured into 3N hydrochloric acid (30 mL) and filtered to give a solid (2.86 g). The filtrate was extracted with dichloromethane, washed with 3N hydrochloric acid and with saturated NaHCO₃, dried over MgSO₄, concentrated in vacuo, combined with the previously collected solid, purified by flash chromatography on silica gel eluting with 1% methanol/dichloromethane, and recrystallized from dichloromethane/isooctane to give a solid (2.44 g, mp 220-222°C) which was the DMF adduct of the desired product. The DMF adduct (2.44 g) was dissolved in dioxane (18 mL) and treated with a 4N solution of hydrochloric acid in dioxane (10 mL). The solution was stirred at room temperature for 7.25 hours, heated to reflux for 42 hours, filtered, and concentrated in vacuo. The residue was dissolved in methylene chloride, washed with water, dried over MgSO₄, and reconcentrated in vacuo to give a brown foam (2.43 g) which was a mixture of the desired product and its adduct with DMF. The mixture was purified by flash chromatography on silica gel eluting with 40% ethyl acetate/hexane to give the desired product as a white solid (0.62 g, 19%): mp 211-213°C; ¹H NMR (acetone-d₆) 300 MHz 8.10 (d, J=8.5 Hz, 2H) 7.76 (d, J=8.5 Hz, 2H) 7.54 (d, J=7.7 Hz, 1H) 7.30 (dd, 1H) 7.07 (dd, 1H) 6.94 (d, J=8.1 Hz, 1H) 6.82 (br s, 2H) 4.12 (s, 2H). High resolution mass Calc'd. for C₁₇H₁₂N₄O₂S₂: 368.0402. Found: 368.0368.

EXAMPLE 23



5 4-[1,4-Dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)[1]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 2-fluoro-3-methoxyphenyl)-3-
10 oxo-propanoic acid.

A solution of 2-fluoro-3-methoxybenzyl alcohol (1.57 g, 9.22 mmol), chloroacetic acid (1.72 g, 18.2 mmol), and ethanol (0.04 mL) in 20 mL of anhydrous tetrahydrofuran was added to a mixture of sodium hydride (2.29 g, 95.2 mmol) in 10 mL of anhydrous tetrahydrofuran dropwise over 10 minutes at 0°C. The cooling bath was removed and the solution warmed to room temperature and then was heated to reflux for 14 hours. The solution was cooled to room temperature, acidified with 3N hydrochloric acid, and extracted with ether. The ethereal phase was washed with brine, dried over anhydrous MgSO₄, filtered, and concentrated *in vacuo* to provide a yellow solid (2.04 g, 100%) that was used directly in the next step: ¹H NMR (CDCl₃/300 MHz) 8.60 (brs, 1H), 7.08-6.93 (m, 3H), 4.72 (d, J=1.4 Hz, 2H), 4.17 (s, 2H), 3.88 (s, 3H); ¹⁹F NMR (CDCl₃/282 MHz) -141.50 (t). Mass spectrum M+Li = 221.

Step 2. Preparation of 7-methoxy-8-fluoro-isochroman-4-one.

A solution of 2-fluoro-3-methoxyphenyl)-3-oxo-propanoic acid from Step 1 (1.96 g, 8.6 mmol) in 4 mL of trifluoroacetic acid and 2 mL of trifluoroacetic anhydride was stirred at room temperature for 1 hour. The solution was concentrated *in vacuo*, the residue dissolved in ether, and the ethereal solution was washed with saturated aqueous NaHCO₃, brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo* to give a solid that was purified by flash chromatography to provide 0.37 g (32%) of the desired ketone. The aqueous phase from the NaHCO₃ was acidified, extracted with ether, washed with brine, dried over anhydrous MgSO₄, filtered, and concentrated *in vacuo* to provide 1.13 g, of recovered 2-fluoro-3-methoxyphenyl)3-oxo-propanoic acid: mp 112-118°C; ¹H NMR (CDCl₃/300 MHz) 7.86 (dd, J=8.66, 1.41 Hz, 1H), 7.00 (apparent t, J=8.26 Hz, 1H), 4.96 (s, 2H), 4.31 (s, 2H), 3.97 (s, 3H); ¹⁹F NMR (CDCl₃/282 MHz) -142.2 (d). Mass spectrum M⁺ = 196.

Step 3. Preparation of 7-methoxy-8-fluoro-3-(trifluoroacetyl)isochroman-4-one.


A solution of 7-methoxy-8-fluoro-isochroman-4-one from Step 2 (370 mg, 1.76 mmol) and ethyl trifluoroacetate (290 mg, 2.04 mmol) in 8 mL of anhydrous tetrahydrofuran was treated with a solution of 25% sodium methoxide in methanol (570 mg, 2.64 mmol). The solution was stirred at room temperature for 16 hours, treated with excess 3N hydrochloric acid, and extracted with ether. The ethereal extract was washed with brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo* to afford 450 mg (88%) of pure 7-methoxy-8-fluoro-3-(trifluoroacetyl)isochroman-4-one which was used

directly in the next step without further purification.

Step 4. Preparation of 4-[6-fluoro-1,4-dihydro-7-methoxy-3-(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

7-Methoxy-8-fluoro-3-(trifluoroacetyl)isochroman-4-one from Step 3 (450 mg, 1.47 mmol) and 4-sulfonamidophenylhydrazine hydrochloride (430 mg, 1.92 mmol) were dissolved in 5 mL of anhydrous ethanol and heated to reflux for 45 minutes. The solution was concentrated *in vacuo* and the residue was dissolved in ethyl acetate. The solution was washed with 3N hydrochloric acid, brine, dried over anhydrous MgSO_4 , filtered, and concentrated *in vacuo*. The residue was crystallized from a mixture of isooctane and ethyl acetate to afford 4-[1,4-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide as a white solid (200 mg, 30%): mp 289.5-291.0°C; ^1H NMR (acetone d_6 /300 MHz) 8.12 (d, $J=8.4$ Hz, 2H), 7.90 (d, $J=8.5$ Hz, 2H), 7.07 (dd, $J=8.7$, 8.5 Hz, 1H), 6.83 (br s, 2H), 6.75 (d, $J=8.7$ Hz, 1H), 5.45 (s, 2H), 3.90 (s, 3H); ^{19}F NMR (acetone d_6 /282 MHz) -62.51 (s), -140.97 (d). High resolution mass spectrum Calc'd. for $\text{C}_{18}\text{H}_{13}\text{F}_4\text{N}_3\text{O}_4\text{S}$: 443.0563. Found: 443.0570.

EXAMPLE 24



4-[3-(Difluoromethyl)-1,5-dihydro-7-methyl-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide

5

Step 1. Preparation of S-(3-methylbenzyl)-isothiuronium chloride.

Thiourea (26.19 g, 344 mmol) was added to a solution of α -chloro-m-xylene (48.21 g, 343 mmol) in methanol (50 mL). The reaction was heated to reflux and additional methanol (10 mL) was added to dissolve all of the thiourea. After 64.3 hours, the reaction was filtered and dried in vacuo to give a white solid (68.15 g, 92%): mp 182-186 °C. ^1H NMR (DMSO- d_6 300 MHz) 9.34 (br s, 4H) 7.22 (m, 3H) 7.12 (m, 1H) 4.48 (s, 2H) 2.27 (s, 3H).

15

Step 2. Preparation of 3-(3-methylphenylthio)propanoic acid.

The thiuronium salt from Step 1 (10.99 g, 51 mmol) was added to sodium chloroacetate (8.86 g, 76 mmol), ethanol (95 mL), and water (10 mL). After heating to reflux, a solution of NaOH (9.05 g, 226 mmol) in water (50 mL) was added to the reaction dropwise over seven minutes. After stirring for 3.6 hours, the reaction was acidified with concentrated hydrochloric acid, extracted with ether, washed with brine, dried over MgSO_4 , and concentrated in vacuo to give a white solid (9.95 g, 100%): mp 73-75.5 °C. ^1H NMR (CDCl_3 300 MHz) 7.16 (m, 4H) 3.83 (s, 2H) 3.12 (s, 2H) 2.35 (s, 3H). Mass spectrum: $M^+=196$.

20

25

30

Step 3. Preparation of 7-methylisothiochroman-4-one.

The acid from Step 2 (6.06 g, 31 mmol) was dissolved in trifluoroacetic acid (11 mL), treated with trifluoroacetic anhydride (5 mL) and stirred at room temperature for 0.33 hour. The reaction was poured into 10% Na_2CO_3 (100 mL) and extracted with ether, washed with brine, dried over MgSO_4 , concentrated in vacuo, and recrystallized from ether/hexane to give 7-

35

chloroisothiochroman-4-one (2.25 g, 41%) as a white solid: mp 79.5-82 °C. ¹H NMR (CDCl₃ 300 MHz) 7.97 (d, J=8.1 Hz, 1H) 7.17 (d, J=8.1 Hz, 1H) 7.00 (s, 1H) 3.87 (s, 2H) 3.52 (s, 2H) 2.37 (s, 3H). Mass spectrum: M+H=179.

5

Step 4. Preparation of 7-methyl-3-(difluoroacetyl)isothiochroman-4-one.

Ethyl difluoroacetate (1.27 g, 10.3 mmol) was dissolved in ether (100 mL). To the stirred solution was added 25 weight % sodium methoxide (2.47 mL, 10.8 mmol) followed by 7-methylisothiochroman-4-one from Step 3 (1.83 g, 10.27 mmol) dissolved in ether (50 mL). The reaction was stirred at room temperature for 16 hours and treated with 1N hydrochloric acid. The organic layer was collected, washed with brine, dried over MgSO₄, concentrated in vacuo. The crude product was used without further purification.

15


Step 5. Preparation of 4-[3-(difluoromethyl)-1,5-Dihydro-7-methyl-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

20

4-Sulfonamidophenylhydrazine hydrochloride (2.67 g, 11.9 mmol) was added to a stirred solution of the diketone from Step 4 (2.35 g, 9.17 mmol) in ethanol (75 mL). The reaction was heated to reflux and stirred for 16 hours. The reaction was concentrated in vacuo and the residue was dissolved in ethyl acetate, washed with water and brine, dried over MgSO₄, concentrated in vacuo, and crystallized from ethyl acetate/isooctane to give the pyrazole as a white solid (1.98 g, 53%): ¹H NMR (acetone-d₆ 300 MHz) 2.34 (s, 3H), 4.06 (s, 2H), 6.8 (m, 3H), 6.96 (t, 1H, J = 54 Hz), 7.0 (m, 1H), 7.34 (s, 1H), 7.78 (d, 2H, J = 8.66 Hz), 8.06 (d, 2H, J = 8.66 Hz). ¹⁹F NMR (acetone-d₆ 282 MHz) -115.5, d, J = 54 Hz.

30
35

EXAMPLE 25



5 4-[7-Chloro-3-(difluoromethyl)-1,5-dihydro-
 [2]benzothiopyrano[4,3-c]pyrazol-1-
 yl]benzenesulfonamide

10 Step 1. Preparation of S-(3-chlorobenzyl)-isothiuronium
 chloride.

 3-Chlorobenzyl chloride (24.2 g, 0.15 mol) and
 thiourea (11.4 g, 0.15 mol) were dissolved in methanol (70
 mL) and heated to reflux for 16 hours. The reaction was
 cooled to room temperature and a precipitate formed. Ether
15 (150 mL) was added to complete the precipitation of
 product. The crystals were isolated by filtration and
 washed with ether (100 mL). After drying in vacuo, 31.9 g
 (90%) of pure S-(3-chlorobenzyl)- isothiuronium chloride
 was obtained: ¹H NMR (CD₃OD) 4.43(s, 2H) 7.36(s, 3H)
20 7.47(s, 1H).

Step 2. Preparation of 3-chlorobenzylmercaptoacetic acid.

 S-(3-Chlorobenzyl)-isothiuronium chloride from Step 1
 (11.86 g, 50 mmol) and chloroacetic acid (7.1 g, 75 mmol)
25 were dissolved in ethanol (100 ml) and heated to 80 °C in a
 4-neck flask equipped with nitrogen inlet, condenser and
 addition funnel. A solution of NaOH (10 g) in H₂O (100 mL)
 and ethanol (50 mL) was added dropwise over 1 hour to this
 hot solution. The reaction was then heated to 100 °C for 4
30 hours. The reaction was cooled to room temperature,
 acidified with concentrated hydrochloric acid (45 mL) and

extracted with ether (500 mL). The organic layer was washed with brine (300 ml), dried over MgSO_4 and concentrated in vacuo to yield 10.84 g (100%) of 3-chlorobenzylmercaptoacetic acid which was used without further purification: ^1H NMR (CDCl_3 300 MHz) 3.08 (s, 2H) 3.80 (s, 2H) 7.23 (m, 3H) 7.34 (s, 1H), 9.07 (broad s, 1H).

Step 3. Preparation of 7-chloroisothiochroman-4-one.

3-Chlorobenzylmercaptoacetic acid (Step 2) (3.35 g) was dissolved in trifluoroacetic acid (50 mL). Trifluoroacetic acid anhydride (25 mL) was added and the reaction was heated at reflux, under nitrogen, for 16 hours. The solution was carefully poured into 10% Na_2CO_3 solution (500 mL) which was stirring vigorously. The organics were extracted into ether (500 mL) and washed with brine (300 mL), dried over MgSO_4 and concentrated in vacuo. The resulting brown solid was purified by SiO_2 chromatography, eluting with a 0-10% gradient of ethyl acetate in hexane to yield 7-chloroisothiochroman-4-one, (1.57 g, 51%): ^1H NMR (CDCl_3 300 MHz) 3.49 (d, 2H, $J = 0.8$ Hz), 3.8 (s, 2H), 7.16 (m, 1H), 7.3 (dd, 1H, $J = 2.0, 8.4$ Hz), 7.96 (d, 1H, $J = 8.5$ Hz).


Step 4. Preparation of 7-chloro-3-(difluoroacetyl) isothiochroman-4-one.

Ethyl difluoroacetate (1.37 g, 11.1 mmol) was dissolved in diethyl ether (100 mL). To the stirred solution was added 25 weight % sodium methoxide (2.7 ml, 11.7 mmol), followed by 7-chloroisothiochroman-4-one (Step 3) (2.2 g, 11.1 mmol) dissolved in diethyl ether (50 mL). The reaction was stirred at room temperature for 16 hours, then treated with 1N hydrochloric acid. The organic layer was collected, washed with brine, dried over MgSO_4 , concentrated in vacuo. The crude product was used without further purification.

Step 5. Preparation of 4-[7-chloro-3-(difluoromethyl)-1,5-dihydro-[2]benzothiopyrano [4,3-c]pyrazol-1-yl]benzenesulfonamide.

A mixture of the product from Step 4 (11.1 mmol) and
5 4-sulphonamidophenylhydrazine hydrochloride (3.2 g, 14.4 mmol) were dissolved in ethanol (100 mL) and heated to reflux for 16 hours. The reaction was concentrated in vacuo and the resulting solid was dissolved in ethyl acetate (200 mL). The organics were washed with water (200
10 mL), brine (150 mL), dried over MgSO₄ and concentrated in vacuo. The resulting oil was chromatographed on SiO₂ eluting with a gradient of ethyl acetate (from 10-50%) in hexane to yield pure tricyclic pyrazole (1.9 g, 40%): ¹H NMR (acetone-d₆ 300 MHz) 4.13 (s, 2H), 6.8 (s, 2H), 6.9 (m,
15 1H), 7.0 (t, 1H, J = 78 Hz), 7.25 (m, 1H), 7.6 (s, 1H), 7.8 (d, 2H, J = 8.66 Hz), 8.1 (d, 2H, J = 8.66 Hz). ¹⁹F NMR (acetone-d₆ 282 MHz) -115.5 (d, J = 78 Hz).

EXAMPLE 26



**4-[1,5-Dihydro-7-methoxy-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide**

Step 1. Preparation of 3-(3-methoxybenzyloxy)acetic acid.

A suspension of sodium hydride (5.49 g, 0.217 mol) in
80 mL of anhydrous THF was cooled to 0 °C and treated with
30 a solution of 3-methoxybenzyl alcohol (10.0 g, 72.4 mmol)
and chloroacetic acid (10.26 g, 0.108 mol) in 80 mL of THF

over ca. 0.25 hour. The solution was stirred at 0 °C for 2 hours and then warmed to room temperature for 14 hours.

The solution was cooled to 0 °C, treated with 50 mL of 6N HCl and extracted with dichloromethane. The

- 5 dichloromethane solution was washed with brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo* to afford 13.48 g (95%) of pure 3-(3-methoxybenzyloxy)acetic acid that was used directly in the next step: ¹H NMR (CDCl₃ 300 MHz) 7.31-7.26 (m, 1H), 6.94-6.85 (m, 3H), 4.63 (s, 2H), 4.15 (s, 2H), 3.82 (s, 3H).
- 10

Step 2. Preparation of 7-methoxyisochroman-4-one.

- A solution of 3-(3-methoxybenzyloxy)acetic acid (Step 1) (6.50 g, 33.1 mmol) in 20 mL of trifluoroacetic acid was
- 15 treated with trifluoroacetic anhydride (5.15 g, 36.4 mmol) and stirred at room temperature for 0.75 hour. The solution was concentrated *in vacuo* and the residue was purified by flash chromatography over silica gel eluting with 10% ethyl acetate in hexane to afford 3.96 g (67%) of
- 20 7-methoxyisochroman-4-one: ¹H NMR (CDCl₃/300 MHz) 8.01 (d, J=8.66 Hz, 1H), 6.90 (dd, J=8.66, 2.42 Hz, 1H), 6.65 (d, J=2.42 Hz, 1H), 4.84 (s, 2H), 4.32 (s, 2H), 3.87 (s, 3H).


Step 3. Preparation of 4-[1,5-dihydro-7-methoxy-3-(trifluoromethyl)-[2]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

25

- A mixture of 7-methoxyisochroman-4-one (3.96 g, 22.2 mmol) and ethyl trifluoroacetate (2.90 mL, 5.8 mmol) was dissolved in 30 mL of diethyl ether and treated with a
- 30 solution of 25% sodium methoxide in methanol (5.8 mL, 26.7 mmol). The solution was stirred at room temperature for 1 hour. The solution was diluted with 1N HCl and extracted with ethyl acetate, washed with brine, dried over anhydrous MgSO₄, filtered and concentrated *in vacuo* to afford a
- 35 solid. The crude reaction product was dissolved in 50 mL of ethanol and treated with 4-sulfonamidophenylhydrazine hydrochloride (5.47 g, 24.4 mmol). This solution was

145

heated to reflux for 3.5 hours, cooled to room temperature, diluted with 100 mL of 1N HCl. After cooling the solution to 0 °C for ca. 0.5 hour, crystals of pure 4-[1,5-dihydro-7-methoxy-3- (trifluoromethyl)-[2]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide formed that were isolated by filtration and dried in vacuo to afford 7.56 g (80%): mp 275 °C (dec). ¹H NMR (CDCl₃ 300 MHz) 7.91 (d, J=8.66 Hz, 2H), 7.54 (d, J=8.66 Hz, 2H), 6.69-6.54 (m, 5H), 5.07 (s, 2H), 3.63 (s, 3H). ¹⁹F NMR (CDCl₃ 282 MHz) -62.01(s). Anal. calc'd. for C₁₈H₁₄F₃N₃O₄S: C, 50.82; H, 3.32; N, 9.88; S, 7.54. Found: C, 50.93; H, 3.38; N, 9.94; S, 7.57.

EXAMPLE 27

4-[1,5-Dihydro-7,8,9-trimethoxy-3-(trifluoromethyl)-[2]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 3-(3,4,5-trimethoxybenzyloxy)acetic acid.

A suspension of sodium hydride (10.2 g, 0.403 mol) in 80 mL of anhydrous THF was cooled to 0 °C and treated with a solution of 3,4,5-trimethoxybenzyl alcohol (8.00 g, 40.4 mmol) and chloroacetic acid (7.63 g, 80.7 mmol) in 80 mL of THF over 1 hour. The solution was stirred at 0 °C for 1 hour and heated to reflux for 14 hours. The solution was cooled to 0 °C, treated with 30 mL of methanol, 10 mL of water, and then the solution was extracted with dichloromethane. The dichloromethane solution was washed

with brine, dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo* to afford 10.4 g, 100% of pure 3-(3-methoxybenzyloxy)acetic acid that was used directly in the next step.

5

Step 2. Preparation of 6,7,8-trimethoxyisochroman-4-one.

A solution of 3-(3,4,5-trimethoxybenzyloxy)acetic acid (Step 1) (10.0 g, 39 mmol) in 30 mL of trifluoroacetic acid was treated with trifluoroacetic anhydride (15 mL) and
10 stirred at room temperature for 20 minutes. The solution was concentrated *in vacuo* and used directly in the next step without purification: ^1H NMR (CDCl_3 300 MHz) 6.56 (s, 2H), 4.58 (s, 2H), 4.16 (s, 2H), 3.86 (s, 6H), 3.83 (s, 3H).

15


Step 3. Preparation of 4-[1,5-dihydro-7,8,9-trimethoxy-3-(trifluoromethyl)-[2]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

The crude product from Step 2 was dissolved in 40 mL
20 of diethyl ether, mixed with ethyl trifluoroacetate (3.35 mL, 28.1 mmol), treated with 25% sodium methoxide in methanol (10 mL, 30.7 mmol) and stirred at room temperature for 16 hours. The solution was diluted with 30 mL of 1N HCl, extracted with ethyl acetate, washed with brine, dried
25 over anhydrous MgSO_4 , filtered and concentrated *in vacuo* to afford 6.3 g of an oil. The oil was dissolved in ethanol, mixed with 4-sulfonamidophenylhydrazine hydrochloride (6.3 g, 28.2 mmol) and the solution was heated to reflux for 16 hours. The solution was cooled to room temperature,
30 diluted with 30 mL of 1N HCl, extracted with ethyl acetate, washed with brine, dried over anhydrous MgSO_4 , filtered and concentrated *in vacuo* to afford a solid that showed two major components by TLC. The material was purified by flash chromatography over silica gel, eluting with 20%
35 ethyl acetate in hexane to afford 877 mg of material that was crystallized from hexane to afford 400 mg (2%) of pure 4-[1,5-dihydro-6,7,8-trimethoxy-3-(trifluoromethyl)-

147

[2]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide from 3-(3,4,5-trimethoxybenzyloxy)acetic acid: mp 130 °C. ^1H NMR (CDCl_3 300 MHz) 7.99 (d, $J=8.66$ Hz, 2H), 7.63 (d, $J=8.66$ Hz, 2H), 6.65 (s, 1H), 5.13 (s, 2H), 4.83 (s, 2H), 3.92 (s, 3H), 3.76 (s, 3H), 3.26 (s, 3H). ^{19}F NMR (CDCl_3 282 MHz) -62.10(s). High resolution mass spectrum calc'd. For $\text{C}_{20}\text{H}_{18}\text{F}_3\text{N}_3\text{O}_6\text{S}$: 485.0885. Found: 485.0779.

EXAMPLE 28



4-[1,5-Dihydro-7-methyl-3-(trifluoromethyl)- [2]benzopyrano[4,3-c]pyrazol-1- yl]benzenesulfonamide

Step 1. Preparation of 3-(3-methylbenzyloxy)acetic acid.

A suspension of sodium hydride (1.96 g, 75.6 mmol) in 80 mL of anhydrous THF was cooled to 0°C and treated with a solution of 3-methylbenzyl alcohol (3.16 g, 25.9 mmol) and chloroacetic acid (3.67 g, 38.8 mmol) in 80 mL of THF over ca. 1 hour. The solution was stirred at 0 °C for 1.5 hours and heated to reflux for 16 hours. The solution was cooled to 0 °C, treated with 50 mL of 6N HCl and extracted with dichloromethane. The dichloromethane solution was washed with brine, dried over anhyd. MgSO_4 , filtered and concentrated in vacuo to afford 2.67 g (57%) of pure 3-(3-methylbenzyloxy)acetic acid that was used directly in the next step. ^1H NMR (CDCl_3 300 MHz) 7.28-7.12 (m, 4H), 4.62 (s, 2H), 4.14 (s, 2H), 2.36 (s, 3H).

Step 2. Preparation of 7-methylisochroman-4-one.

A solution of 3-(3-methylbenzyloxy)acetic acid (Step 1) (2.67 g, 14.8 mmol) in 25 mL of trifluoroacetic acid was treated with trifluoroacetic anhydride (2.5 mL, 17.8 mmol) and stirred at room temperature for 16 hours. The solution was concentrated in vacuo and the residue was dissolved in dichloromethane, washed with sat. aq. NaHCO₃, brine, dried over anhyd. MgSO₄, filtered and concentrated in vacuo to afford an oil, 1.45 g, 60%. Examination of the NMR spectrum revealed that the product was a 3:1 mixture of 7-methylisochroman-4-one:5-methylisochroman-4-one. Flash chromatography of the mixture eluting with 10% ethyl acetate in hexane provided the desired isomer 7-methylisochroman-4-one [¹H NMR (CDCl₃ 300 MHz) 7.94 (d, J=7.86 Hz, 1H), 7.22 (d, J=7.86 Hz, 1H), 7.02 (s, 1H), 4.85 (s, 2H), 4.35 (s, 2H), 2.42 (s, 3H)].


Step 3. Preparation of 4-[1,5-dihydro-7-methyl-3-(trifluoromethyl)-[2]benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

The product from Step 2 was dissolved in 20 mL of diethyl ether, mixed with ethyl trifluoroacetate (0.12 mL, 9.5 mmol), treated with 25% sodium methoxide in methanol (2.2 mL, 10.4 mmol) and stirred at room temperature for 16 hours. The solution was diluted with 10 mL of 1N HCl, extracted with ethyl acetate, washed with brine, dried over anhydrous MgSO₄, filtered and concentrated in vacuo to afford an oil. The oil was dissolved in ethanol, mixed with 4-sulfonamidophenylhydrazine hydrochloride (2.12 g, 9.5 mmol) and the solution was heated to reflux for 3 hours. The solution was cooled to room temperature and diluted with 30 mL of 0.5N HCl. The solution was chilled in an ice bath, whereupon crystals formed that were isolated by filtration and dried in vacuo to afford 1.27 g, 36% of pure product: mp 285 °C (dec). ¹H NMR (CDCl₃ 300 MHz) 8.05 (d, J=8.66 Hz, 2H), 7.70 (d, J=8.66 Hz, 2H), 7.05 (s, 1H), 6.97 (d, J=7.96 Hz, 1H), 6.77 (d, J=7.96, 1H), 5.20 (s, 2H), 2.31 (s, 3H). ¹⁹F NMR (CDCl₃ 282 MHz)

-62.11(s). High resolution mass spectrum calc'd. for $C_{18}H_{15}F_3N_3O_3S$ (MH^+): 410.0786. Found: 410.0754.

EXAMPLE 29

5



10

4-[6,8-Difluoro-1,5-dihydro-7-methoxy-3-(trifluoromethyl)[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

Step 1. Preparation of 2,6-difluoroanisole.

A solution of 2,6-difluorophenol (30.00 g, 230 mmol) in 2.5 N sodium hydroxide (95 mL, 238 mmol) was treated with dimethylsulfate (33.32 g, 264 mmol) and stirred for 3.3 hours at 50 °C. Additional dimethylsulfate (13.33 g, 105 mmol) was added and the reaction was stirred another 0.6 hour. The reaction mixture was extracted with ethyl acetate, washed with 2.5 N sodium hydroxide, brine, dried over $MgSO_4$, and concentrated in vacuo to give a clear oil (25.93 g, 78%): 1H NMR ($CDCl_3$ 300 MHz) 6.87 (m, 3H) 3.98 (s, 3H); ^{19}F NMR ($CDCl_3$ 282 MHz) -129.48 (t). Mass spectrum: $M^+=144$.

Step 2. Preparation of 2,4-difluoro-3-methoxybenzoic acid.

A solution of potassium tert-butoxide (22.74 g, 203 mmol) in anhydrous tetrahydrofuran (235 mL) was cooled to -78 °C and treated with a 1.6 M solution of n-butyllithium (120 mL, 192 mmol) in hexanes. After stirring for 30 minutes, 2,6-difluoroanisole (Step 1) (25.89 g, 180 mmol) was added and the reaction was stirred an additional 7.4

hours. The reaction was poured into dry ice and warmed to room temperature. Water (250 mL) was added, and after extracting with ether (160 mL), the aqueous layer was acidified with concentrated HCl, and filtered to give a
5 yellow solid (3.71 g, 11%): mp 176-182 °C. ¹H NMR (CDCl₃ 300 MHz) 7.70 (m, 1H) 6.98 (m, 1H) 4.01 (s, 3H); ¹⁹F NMR (CDCl₃ 282 MHz) -119.06 (m) -123.33 (m). Mass spectrum: M+Li=195.

10 Step 3. Preparation of 2,4-difluoro-3-methoxybenzyl alcohol.

A solution of 2,4-difluoro-3-methoxybenzoic acid (Step 2) (3.65 g, 19 mmol) in anhydrous tetrahydrofuran (24 mL) was cooled in an ice bath, and treated with borane dimethyl
15 sulfide complex (4 mL, 40 mmol). The reaction was stirred at room temperature for 17 hours, quenched by the slow addition of methanol, and concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with NaHCO₃, brine, dried over MgSO₄, and concentrated in vacuo to give
20 a brown oil (2.93 g, 87%): ¹H NMR (CDCl₃ 300 MHz) 7.01 (m, 1H) 6.86 (m, 1H) 4.66 (s, 2H) 3.97 (s, 3H); ¹⁹F NMR (CDCl₃ 282 MHz) -129.67 (m) -135.09 (m). Mass spectrum: M+=174.

25 Step 4. Preparation of 2,4-difluoro-3-methoxybenzyl chloride.

A solution of 2,4-difluoro-3-methoxybenzyl alcohol (Step 3) (2.93 g, 17 mmol) in concentrated HCl (15 mL) was treated with HCl gas for 3 minutes. The reaction was stirred at room temperature (21 hours). The reaction
30 mixture was extracted with ether, dried over MgSO₄, and concentrated in vacuo to give a brown oil (2.30 g, 71%): ¹H NMR (CDCl₃ 300 MHz) 7.03 (m, 1H) 6.89 (m, 1H) 4.58 (s, 2H) 4.01 (s, 3H); ¹⁹F NMR (CDCl₃ 282 MHz) -127.88 (m) -132.78 (m). Mass spectrum: M+=192.

35

Step 5. Preparation of S-(2,4-difluoro-3-methoxybenzyl)-isothiuronium chloride.

Thiourea (0.90 g, 12 mmol) was added to a solution of 2,4-difluoro-3-methoxybenzyl chloride (Step 4) (2.30 g, 12 mmol) in methanol (7 mL). The reaction was heated to reflux for 2.9 hours and concentrated *in vacuo* to give a white solid (3.18 g, 100%): mp 144-151 °C. ¹H NMR (DMSO-d₆ 300 MHz) 9.35 (br. d, 3H) 7.26 (m, 1H) 7.15 (m, 1H) 4.57 (s, 2H) 3.91 (s, 3H); ¹⁹F NMR (DMSO-d₆ 282 MHz) -128.73 (m) -131.41 (m).

10 Step 6. Preparation of 3-(2,4-difluoro-3-methoxyphenylthio)propanoic acid.

A 100 mL flask was charged with the thiouronium salt from Step 5 (3.18 g, 12 mmol), sodium chloroacetate (2.36 g, 20 mmol), ethanol (10 mL), and water (10 mL). After heating to reflux, a solution of NaOH (2.51 g, 63 mmol) in water (10 mL) was added to the reaction dropwise. After stirring for 15.5 hours, the reaction was acidified with concentrated HCl, extracted with ether, washed with brine, dried over MgSO₄, concentrated *in vacuo*, and recrystallized from ether/hexane to give a brown oil (2.33 g, 70%): ¹H NMR (CDCl₃ 300 MHz) 8.4 (br. s, 1H) 6.98 (m, 1H) 6.85 (m, 1H) 3.99 (s, 3H) 3.85 (s, 2H) 3.17 (s, 2H); ¹⁹F NMR (CDCl₃ 282 MHz) -129.33 (m) -132.16 (m). Mass spectrum: M+Li=255.

25 Step 7. Preparation of 6,8-difluoro-7-methoxyisothiochroman-4-one.

The acid from Step 6 (2.30 g, 9.3 mmol) was dissolved in trifluoroacetic acid (10 mL), treated with trifluoroacetic anhydride (5 mL) and stirred at room temperature for 1.3 hours. The reaction was poured into 10% Na₂CO₃ (150 mL) and extracted with ethyl acetate, washed with brine, dried over MgSO₄, and concentrated *in vacuo* to give a brown solid (0.45 g) which was used in the next step without further purification.

35

Step 8. Preparation of 6,8-difluoro-7-methoxy-3-(trifluoroacetyl)isothiochroman-4-one.


Ethyl trifluoroacetate (0.30 g, 2.1 mmol) was added to a solution of the isothiochromanone from Step 7 (0.45 g, 2.0 mmol) in ether (5 mL). The reaction was treated with 25 weight % NaOMe (0.52 g, 2.4 mmol) and stirred at room temperature for 6.5 hours, then treated with 3N HCl. The organic layer was collected, washed with brine, dried over MgSO₄, and concentrated in vacuo to give a brown oil (0.43 g) which was used in the next step without further purification.

10

Step 9. Preparation of 4-[6,8-difluoro-1,5-dihydro-7-methoxy-3-(trifluoromethyl)[2]benzothioapyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

The 4-sulfonamidophenylhydrazine hydrochloride (0.38 g, 1.7 mmol) was added to a stirred solution of the diketone (Step 8) (0.43 g, 1.3 mmol) in ethanol (10 mL). The reaction was heated to reflux and stirred for 16.6 hours. The reaction mixture was filtered, and the filtrate was concentrated in vacuo, dissolved in ethyl acetate, washed with water and brine, dried over MgSO₄, concentrated in vacuo, and passed through a column of silica gel to give the pyrazole as a brown oil (0.23 g, 37%): ¹H NMR (acetone-d₆ 300 MHz) 8.11 (d, J=8.7 Hz, 2H) 7.87 (d, J=8.7 Hz, 2H) 6.84 (br s, 2H) 6.57 (d, J=11.9 Hz, 1H) 4.20 (s, 2H) 4.03 (t, J=1.0 Hz, 3H); ¹⁹F NMR (acetone-d₆ 282 MHz) -63.33 (s) -130.85 (m) -132.61 (m). High resolution mass spectrum calc'd. for C₁₈H₁₂F₅N₃O₃S₂: 477.0240. Found: 477.0205.

25

EXAMPLE 30

5 **4-[3-(Difluoromethyl)-1,5-dihydro-7-**
 methyl[2]benzopyrano[4,3-c]pyrazol-1-
 yl]benzenesulfonamide

10 Step 1. Preparation of (3-methylphenyl)-3-oxo-propanoic
 acid.

 A solution of 3-methylbenzyl alcohol (4.93 g, 40 mmol), chloroacetic acid (7.68 g, 81 mmol), and ethanol (0.04 mL) in 100 mL of anhydrous tetrahydrofuran was added to a mixture of sodium hydride (16.12 g, 403 mmol) in 50 mL
15 anhydrous tetrahydrofuran dropwise over 55 minutes at 0 °C. The cooling bath was removed and the solution was heated to reflux for 14.25 hours. The solution was cooled to room temperature, quenched with water and extracted with ether. The aqueous layer was acidified, extracted with ether,
20 washed with brine, dried over MgSO₄, and concentrated in vacuo to give (3-methylphenyl)-3-oxo-propanoic acid as an orange oil (6.58 g) which also contained chloroacetic acid. The oil was used in the next step without purification.

25 Step 2. Preparation of 7-methylisochroman-4-one.

 The acid from Step 1 (6.58 g, 36 mmol) was dissolved in trifluoroacetic acid (20 mL), treated with trifluoroacetic anhydride (11 mL) and stirred at room temperature for 14.75 hours. The reaction was poured into
30 10% Na₂CO₃ (150 mL) and extracted with ether, washed with brine, dried over MgSO₄, concentrated in vacuo, and passed


through a column of silica gel with 15% ethyl acetate/hexane to give a yellow solid (1.09 g, 18%): ¹H NMR (CDCl₃ 300 MHz) 7.95 (d, J=7.9 Hz, 1H) 7.23 (d, J=7.9 Hz, 1H) 7.02 (s, 1H) 4.85 (s, 2H) 4.34 (s, 2H) 2.42 (s, 3H).

Step 3. Preparation of 3-(difluoroacetyl)-7-methylisochroman-4-one.

Ethyl difluoroacetate (1.02 g, 8.2 mmol) was added to a solution of the isochromanone from Step 2 (1.07 g, 6.6 mmol) in ether (20 mL). The reaction was treated with 25 weight % NaOMe (1.89 g, 8.7 mmol) and stirred at room temperature for 3.1 hours. The reaction mixture was filtered to give a yellow solid which was dissolved in 3N HCl, extracted with ether, washed with brine, dried over MgSO₄, and concentrated in vacuo to give a yellow solid (0.79 g) which was used in the next step without further purification.

Step 4. Preparation of 4-[3-(difluoromethyl)-1,5-dihydro-7-methyl[2]benzopyranof[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-Sulfonamidophenylhydrazine hydrochloride (0.77 g, 3.4 mmol) was added to a stirred solution of the diketone from Step 3 (0.75 g, 3.1 mmol) in ethanol (10 mL). The reaction was heated to reflux and stirred for 2.25 hours. The reaction mixture was filtered while hot and recrystallized from ethyl acetate/hexane to give a yellow solid (0.41 g, 34%): mp 178-179 °C (dec). ¹H NMR (DMSO-d₆ 300 MHz) 8.00 (d, J=8.7 Hz, 2H) 7.84 (d, J=8.5 Hz, 2H) 7.56 (br s, 2H) 7.25 (s, 1H) 7.12 (t, J=53.6 Hz, 1H) 7.08 (d, J=8.1 Hz, 1H) 6.72 (d, 8.1 Hz) 5.28 (s, 2H) 2.29 (s, 3H); ¹⁹F NMR (DMSO-d₆ 282 MHz) -114.42 (d). High resolution mass spectrum calc'd. for C₁₈H₁₅F₂N₃O₃S: 391.0802. Found: 391.0798.

EXAMPLE 31

5 **4-[3-(Difluoromethyl)-1,5-dihydro-7-fluoro-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide**

10 **Step 1. Preparation of S-(3-fluorobenzyl)-isothiouronium**
chloride.

A solution of 3-fluorobenzyl chloride (49.95 g, 0.346 mol) dissolved in 60 mL of anhydrous methanol was treated portion wise with thiourea (26.30 g, 0.346 mol) over a period of 0.25 hour at room temperature. The solution was warmed to reflux for 0.5 hour, cooled to 5 °C and the product was isolated by filtration to afford 59.17 g. Concentration of the filtrate and chilling in an ice bath afforded an additional 9.93 g of pure S-(3-fluorobenzyl)isothiouronium chloride (96%): mp 201-203 °C.

20 **Step 2. Preparation of 3-(3-fluorophenylthio)propanoic acid.** S-(3-Fluorobenzyl)-isothiouronium chloride from step 1 (25.0 g, 94 mmol) was added to sodium chloroacetate (16.4 g, 141 mmol), sodium hydroxide pellets (15.0 g, 376

25 mmol), ethanol (150 mL), and water (100 mL). This mixture was stirred for 16 hours at room temperature, then concentrated hydrochloric acid was added to pH 1. The solution was extracted with ether (2 X 100 mL), combined and washed with brine, dried over MgSO₄, and concentrated.
30 A brown solid was recrystallized from ether/hexanes (18.7 g, 95%): mp 86-88 °C.

Step 3. Preparation of 7-fluoroisothiochroman-4-one.

3-(3-Fluorophenylthio)propanoic acid from step 2 (18.0 g, 90 mmol) was dissolved in trifluoroacetic acid (30 mL), treated with trifluoroacetic anhydride (90 mL) and heated to reflux for 16 hours. Ether (200 mL) was added, and the mixture was washed with saturated aqueous NaHCO₃ solution (2 X 150 mL) and water. After drying over MgSO₄, the solution was concentrated to a brown solid (13.0 g, 79%). This material was used in the next step without further purification or characterization.

Step 4. Preparation of 3-difluoroacetyl-7-fluoroisothiochroman-4-one.

7-Fluoroisothiochroman-4-one from step 3 (2.0 g, 11 mmol) was dissolved in anhydrous ethyl ether (50 mL). Sodium methoxide 25% in methanol (2.61 g, 12 mmol) and ethyl difluoroacetate (1.49 g, 12 mmol) were added. After stirring for 16 hours at room temperature the mixture was washed with 1 N hydrochloric acid, brine, and water, dried over MgSO₄ and concentrated. This crude material was used in the next step without further purification or characterization.


Step 5. Preparation of 4-[3-(difluoromethyl)-1,5-dihydro-7-fluoro-2-benzothiopyran[4,3-c]pyrazol-1-yl]benzenesulfonamide.

7-Fluoro-3-(difluoroacetyl)isothiochroman-4-one from Step 4 (2.9 g, 11 mmol) was dissolved in ethanol (100 mL), and 4-sulfonamidophenylhydrazine hydrochloride (2.68 g, 12 mmol) was added. The mixture was heated to reflux for 16 hours. After cooling, water was added until crystals appeared. A dark brown solid was collected by filtration (1.7 g, 37%): mp 260-262 °C. ¹H NMR (DMSO-d₆ 300 MHz) 7.96 (d, J=8.4 Hz, 2H), 7.75 (d, J=8.4 Hz, 2H), 7.54 (s, 2H), 7.42 (dd, J=2.7, 9.3 Hz, 1H), 7.17 (t, J=53.7 Hz, 1H), 7.09 (dt, J=2.4, 8.4 Hz, 1H), 6.77 (dd, J=5.4, 8.7 Hz, 1H), 4.15

(s, 2H). Anal. calc'd. for $C_{17}H_{12}N_3S_2O_2F_3$: C, 49.63; H, 2.94; N, 10.21. Found: C, 49.55; H, 2.95; N, 10.14.

EXAMPLE 32

5



10

1,5-Dihydro-7-fluoro-1-[(4-methylsulfonyl)phenyl]-
3-(trifluoromethyl)-
[2]benzothiopyrano[4,3,c]pyrazole

Step 1. Preparation of 7-fluoro-3-trifluoroacetyl)isothiochroman-4-one.

7-Fluoroisothiochroman-4-one (Example 31, step 3)
15 (1.82 g, 10 mmol) was dissolved in anhydrous ethyl ether
(50 mL). Sodium methoxide 25% in methanol (2.38 g, 11
mmol) and ethyl trifluoroacetate (1.56 g, 11 mmol) were
added. After stirring for 16 hours at room temperature the
mixture was washed with 1 N hydrochloric acid, brine, and
20 water, dried over $MgSO_4$ and concentrated. This crude
material was used in the next step without further
purification or characterization.


Step 2. Preparation of 1,5-dihydro-7-fluoro-1-[(4-methylsulfonyl)phenyl]-3-(trifluoromethyl)-[2]benzothiopyrano[4,3,c]pyrazole.

7-Fluoro-3-trifluoroacetyl)isothiochroman-4-one (2.9
g, 11 mmol) from Step 1 was dissolved in ethanol (100 mL)
and p-methanesulfonylphenylhydrazine hydrochloride (2.45 g,
30 11 mmol) was added. The mixture was heated to reflux for
16 hours, and concentrated. The resultant solid was
dissolved in ethyl acetate and washed with brine and water,

158

dried over MgSO_4 and concentrated. A dark brown solid was recrystallized from ethyl acetate/hexanes (1.4 g, 32%): mp 193-195 °C. ^1H NMR ($\text{DMSO}-d_6$ 300 MHz) 8.10 (d, $J=8.7$ Hz, 2H), 7.86 (d, $J=8.7$ Hz, 2H), 7.44 (dd, $J=2.7, 9.3$ Hz, 1H), 7.09 (dt, $J=2.7, 8.7$ Hz, 1H), 6.79 (dd, $J=5.7, 8.7$ Hz, 1H), 4.2 (s, 2H), 3.27 (s, 3H). Anal. calc'd. for $\text{C}_{18}\text{H}_{12}\text{N}_2\text{S}_2\text{O}_2\text{F}_4$: C, 50.46; H, 2.82; N, 6.54. Found: C, 50.54; H, 2.84; N, 6.57.

10

EXAMPLE 33

15 **4-[1,5-Dihydro-6,7-methylenedioxy-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide**

Step 1. Preparation of 2,3-methylenedioxybenzoic acid.

20 1,3-Benzodioxole (12.2 g, 100 mmol) was dissolved in tetrahydrofuran (150 mL). This mixture was cooled to -78 °C under nitrogen with stirring while n-butyllithium 1.6 M in hexane (69 mL, 110 mmol) was added dropwise maintaining the temperature below -50 °C. Potassium t-butoxide (12.34 g, 110 mmol) was added and the mixture stirred at -78 °C for 0.25 hour when dry solid carbon dioxide was added. The cooling bath was removed and the mixture was stirred for an additional hour, when it was poured into 250 mL of 1 N hydrochloric acid. The mixture was extracted (3 X 100 mL) with ethyl acetate, combined and the organic portions were 30 extracted into 2.5 N sodium hydroxide solution (2 X 50 mL).

The combined basic aqueous layers were acidified to pH 3 with 1 N hydrochloric acid and extracted with ethyl acetate (3 X 50 mL). The combined organic layers were washed with water, dried over MgSO_4 and concentrated. A light brown solid was recrystallized from ethyl acetate/hexanes (2.9 g, 17%): mp 224-227 °C. ^1H NMR ($\text{DMSO}-d_6$ 300 MHz) 12.9 (bs, 1H), 7.26 (dd, $J=1.2, 8.1$ Hz, 1H), 7.10 (dd, $J=0.9, 7.5$ Hz, 1H), 6.87 (dd, $J=8.1, 7.8$ Hz, 1H), 6.10 (s, 2H). Anal. calc'd for $\text{C}_8\text{H}_6\text{O}_4$: C, 57.84; H, 3.64. Found: C, 57.70; H, 3.73.

Step 2. Preparation of 2,3-methylenedioxybenzyl alcohol.

2,3-Methylenedioxybenzoic acid from Step 1 (2.8 g, 17 mmol) was dissolved in tetrahydrofuran (100 mL) and borane-dimethyl sulfide complex (10 M) was added (5.0 mL, 50 mmol). This mixture was stirred for 16 hours when methanol was added dropwise to destroy unreacted borane. Ethyl acetate (100 mL) was added and the mixture was washed with saturated aqueous NaHCO_3 solution and water twice, dried over MgSO_4 , and concentrated to 2.4 g of a light brown solid. This material was used in the next step without further purification or characterization.

Step 3. Preparation of S-(2,3-methylenedioxybenzyl)-isothiuronium chloride.

A solution of 2,3-methylenedioxybenzyl alcohol from Step 2 (2.4 g, 16 mmol) was dissolved in tetrahydrofuran (25 mL). Triethylamine (2.43 g, 24 mmol) was added followed by the dropwise addition of methanesulfonyl chloride (2.27 g, 20 mmol). After stirring at room temperature for 4 hours, thiourea (1.3 g, 16 mmol) and methanol (50 mL) were added. The mixture was heated to reflux for 16 hours, concentrated and used in the next step without further purification or characterization.

Step 4. Preparation of 3-(2,3-methylenedioxyphenylthio)propanoic acid.

S-(2,3-Methylenedioxybenzyl)-isothiuronium chloride from Step 3 (3.4 g, 16 mmol), sodium chloroacetate (2.8 g, 24 mmol), sodium hydroxide pellets (2.6 g, 64 mmol), ethanol (25 mL), and water (40 mL) were combined and
5 stirred for 16 hours at room temperature. Concentrated hydrochloric acid was added to pH 1, and the solution was extracted with ether (4 X 50 mL). The combined ethereal layers were washed with 1 N hydrochloric acid, dried over MgSO₄, and concentrated (3.62 g, 95%). This mixture was
10 used in the next step without further purification or characterization.

Step 5. Preparation of 6,7-methylenedioxyisothiochroman-4-one.

3-(2,3-Methylenedioxyphenylthio)propanoic acid
15 from Step 4 (3.62 g, 16 mmol) was dissolved in trifluoroacetic acid (5 mL), treated with trifluoroacetic anhydride (15 mL) and stirred at room temperature for 16 hours. After neutralizing with saturated aqueous NaHCO₃, the solution was extracted with ethyl acetate (3 X 50 mL).
20 The combined organic extracts were washed with brine and concentrated. Purification was performed by flash column chromatography eluting with (4:1) hexanes:ethyl acetate. The appropriate fractions were concentrated and recrystallized from ethyl acetate/hexanes to yield a brown
25 solid (0.5 g, 15%): mp 114-116 °C. ¹H NMR (DMSO-d₆ 300 MHz) 7.56 (d, J=8.4 Hz, 1H), 6.95 (d, J=8.1 Hz, 1H), 6.16 (s, 2H), 3.87(s, 2H), 3.57(s, 2H). Anal. calc'd. for C₁₀H₈O₃S: C, 57.68; H, 3.87. Found: C, 57.59; H, 3.93

Step 6. Preparation of 6,7-methylenedioxy-3-(trifluoroacetyl) isothiochroman-4-one.

6,7-Methylenedioxyisothiochroman-4-one from Step 5 (0.44 g, 2.1 mmol) was dissolved in anhydrous ethyl ether (50 mL). Sodium methoxide 25% in methanol (0.54 g, 2.5
35 mmol) and ethyl trifluoroacetate (0.36 g, 2.5 mmol) were added. After stirring for 16 hours at room temperature the mixture was washed with 1 N hydrochloric acid, brine, and

161


water, dried over MgSO_4 and concentrated. This crude material was used in the next step without further purification or characterization.

5 Step 7. Preparation of 4-[1,5-dihydro-7-fluoro-3-(difluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

6,7-Methylenedioxy-3-(trifluoroacetyl)isothiochroman-4-one from Step 6 (0.64 g, 2.1 mmol) was dissolved in ethanol (100 mL) and 4-sulfonamidophenylhydrazine hydrochloride (0.56 g, 2.5 mmol) was added. The mixture was heated to reflux for 16 hours, and concentrated. The resultant oily solid was dissolved in ethyl acetate and washed with brine and water, dried over MgSO_4 and concentrated. Purification was achieved by flash column chromatography eluting with (1:1) ethyl acetate:hexanes. The appropriate fractions were concentrated and recrystallized from ethyl acetate/hexanes. The product was a light brown solid (0.2 g, 21%): mp 272-274 °C. ^1H NMR (DMSO- d_6 300 MHz) 7.98 (d, $J=8.4$ Hz, 2H), 7.78 (d, $J=8.7$ Hz, 2H), 7.44 (bs, 2H), 6.79 (d, $J=8.4$ Hz, 1H), 6.28 (d, $J=8.1$ Hz, 1H), 6.14 (s, 2H), 4.09 (s, 2H). Anal. calc'd for $\text{C}_{18}\text{H}_{12}\text{N}_3\text{S}_2\text{O}_4\text{F}_3$: C, 47.47; H, 2.66; N, 9.23. Found: C, 47.56; H, 2.73; N, 9.16.

25

EXAMPLE 34



30

4-(3-Cyano-1,5-dihydro-7-fluoro-
[2]benzothiopyrano[4,3-

**clpyrazol-1-yl)-N-[(dimethylamino)methylene]-
benzenesulfonamide**

**Step 1. Preparation of 3,4-dihydro- α ,4-dioxo-7-fluoro-1H-
2-benzothiopyran-3-acetic acid, methyl ester.**

7-Fluoroisothiochroman-4-one (Example 31, step 3) (2.0 g, 11 mmol) was dissolved in anhydrous ethyl ether (50 mL), and sodium methoxide 25% in methanol (2.6 g, 12 mmol), and dimethyl oxalate (1.42 g, 12 mmol) were added. After stirring for 16 hours at room temperature, the mixture was washed with 1 N hydrochloric acid, brine, and water, dried over MgSO_4 , concentrated. A brown solid was recrystallized from chloroform/hexanes (2.5 g, 85%): mp 133-135 °C. ^1H NMR ($\text{DMSO}-d_6$ 300 MHz) 15.4 (s, 1H), 8.03 (dd, $J=5.7$, 8.7 Hz, 1H), 7.11 (dt, $J=2.4$, 8.4 Hz, 1H), 6.94 (dd, $J=2.7$, 8.7 Hz, 1H), 3.95 (s, 3H), 3.78 (s, 2H). Anal. calc'd. for $\text{C}_{12}\text{H}_9\text{N}_3\text{SO}_4\text{F}$: C, 53.73; H, 3.38. Found: C, 53.56; H, 3.42.

Step 2. Preparation of 4-[3-(carbomethoxy)-1,5-dihydro-7-fluoro-[2]benzothiopyrano[4,3-clpyrazol-1-yl]benzenesulfonamide.

3,4-Dihydro- α ,4-dioxo-7-fluoro-1H-2-benzothiopyran-3-acetic acid, methyl ester from Step 1 (2.5 g, 9 mmol) was dissolved in methanol (100 mL) and p-sulfonamidophenylhydrazine hydrochloride (2.23 g, 10 mmol) was added. The mixture was heated to reflux for 16 hours. After cooling, water was added until crystals appeared. The product was collected by filtration and recrystallized from methanol/water to yield a dark brown solid (3.6 g, 95%): mp 241-244 °C. ^1H NMR ($\text{DMSO}-d_6$ 300 MHz) 7.99 (d, $J=8.4$ Hz, 2H), 7.8 (d, $J=8.4$ Hz, 2H), 7.56 (s, 2H), 7.42 (dd, $J=2.4$, 9.0 Hz, 1H), 7.08 (dt, $J=2.7$, 8.7 Hz, 1H), 6.75 (dd, $J=5.4$, 8.7 Hz, 1H), 4.18 (s, 2H), 3.85 (s, 3H). Anal. calc'd. for $\text{C}_{18}\text{H}_{14}\text{N}_3\text{O}_4\text{S}_2\text{F}_1$: C, 51.54; H, 3.36; N, 10.02. Found: C, 51.38; H, 3.44; N, 9.94.

Step 3. Preparation of 4-[3-(carboxamido)-1,5-dihydro-7-fluoro-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide.

4-[3-(Carbomethoxy)-1,5-dihydro-7-fluoro-
5 [2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide
from Step 2 (1.0 g, 2.4 mmol), methanol (50 mL), and sodium
cyanide (0.5 g) were placed in a Parr bottle and
pressurized to 20 psig with ammonia gas. After stirring
for 64 hours at room temperature, the mixture was
10 concentrated, dissolved in ethyl acetate, and washed with 1
N hydrochloric acid, and water. The solution was then
dried over MgSO₄ and concentrated. A white solid was
recrystallized from ethyl acetate/hexanes (0.6 g, 62%): mp
297-298 °C. ¹H NMR (DMSO-d₆ 300 MHz) 7.97 (d, J=8.7 Hz,
15 2H), 7.78 (d, J=8.7 Hz, 2H), 7.74 (bs, 1H), 7.53 (s, 2H),
7.51 (bs, 1H), 7.39 (dd, J=2.7, 9.3 Hz, 1H), 7.07 (dt,
J=2.7, 9.0 Hz, 1H), 6.75 (dd, J=5.4, 8.7 Hz, 1H), 4.1 (s,
2H). Anal. calc'd. for C₁₇H₁₃N₄O₃S₂F: C, 50.49; H, 3.24;
N, 13.85. Found: C, 50.67; H, 3.35; N, 13.60.

20


Step 4. Preparation of 4-(3-cyano-1,5-dihydro-7-fluoro-[2]benzothiopyrano[4,3-c]pyrazol-1-yl)-N-[(dimethylamino)methylene]benzenesulfonamide.

N,N'-Dimethylformamide (3 mL) was cooled with stirring
25 to 0 °C and oxalyl chloride (0.7 g, 5.5 mmol) was added
dropwise. 4-[3-(Carboxamido)-1,5-dihydro-7-fluoro-
[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide
from Step 3 (1.0 g, 2.5 mmol) was dissolved in 2 mL of
N,N'-dimethylformamide, added dropwise. The mixture was
30 warmed to room temperature over 0.5 hour when pyridine (2
mL) was added followed by ethyl acetate (100 mL). After
washing with 1 N hydrochloric acid, brine and water, the
mixture was dried over MgSO₄ and concentrated. A light
brown solid was recrystallized from ethyl acetate/hexanes
35 (0.8 g, 73%): mp 232-235 °C. ¹H NMR (DMSO-d₆ 300 MHz) 8.26
(s, 1H), 7.95 (d, J=8.7 Hz, 2H), 7.76 (d, J=8.7 Hz, 2H),
7.45 (dd, J=2.7, 9.3 Hz, 1H), 7.13 (dt, J=2.4, 8.4 Hz, 1H),

164

6.75 (dd, $J=5.4, 8.7$ Hz, 1H), 4.22 (s, 2H), 3.16 (s, 3H), 2.94 (s, 3H). Anal. calc'd. for $C_{20}H_{16}N_5O_2S_2F + 0.5 H_2O$: C, 53.32; H, 3.80; N, 15.54. Found: C, 53.33; H, 3.78; N, 15.50.

5

EXAMPLE 35

10


4-[3-Cyano-1,5-dihydro-7-fluoro-
[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide

4-(1,5-Dihydro-7-fluoro-3-cyano-
[2]benzothiopyrano[4,3-c]pyrazol-1-yl)-N-
[(dimethylamino)methylene]benzenesulfonamide (Example 34)
(0.7 g, 1.6 mmol) was dissolved in tetrahydrofuran (50 mL)
and treated with sodium hydroxide 2.5 N (10 mL). After
stirring for 5 minutes, concentrated hydrochloric acid was
added to pH 1. The mixture was extracted with ethyl
acetate (2 X 30 mL), combined, washed with water, dried
over $MgSO_4$, and concentrated. A tan solid was
recrystallized from ethanol/water (0.6 g, 97%): mp 256-257
 $^{\circ}C$. 1H NMR (DMSO- d_6 300 MHz) 8.00 (d, $J=8.7$ Hz, 2H), 7.81
(d, $J=8.7$ Hz, 2H), 7.58 (s, 2H), 7.46 (dd, $J=2.7, 9.3$ Hz,
1H), 7.11 (dt, $J=2.4, 8.4$ Hz, 1H), 6.77 (dd, $J=5.4, 8.7$ Hz,
1H), 4.22 (s, 2H). Anal. calc'd. for $C_{17}H_{11}N_4O_2S_2F$: C,
52.84; H, 2.87; N, 14.50. Found: C, 52.69; H, 2.95; N,
14.60.

30

EXAMPLE 36

165



5 4-[1,5-Dihydro-7-fluoro-3-(hydroxymethyl)-
 [2]benzothiopyrano[4,3-c]pyrazol-1-
 yl]benzenesulfonamide

4-[3-(Carbomethoxy)-1,5-dihydro-7-fluoro-
[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide
10 (Example 34, Step 2) (1.0 g, 2.4 mmol) was dissolved in
tetrahydrofuran (50 mL) with stirring at room temperature.
Lithium aluminum hydride 1.0 M in tetrahydrofuran (5 mL)
was added dropwise. After 0.5 hour, 1.25 N sodium
hydroxide (10 mL) was added. The mixture was filtered,
15 concentrated, and purified by flash column chromatography
eluting with ethyl acetate : hexanes (2:1). The
appropriate fractions were concentrated and recrystallized
from ethyl acetate/hexanes. Product was a white solid (0.2
g, 21%): mp 186-190 °C. ¹H NMR (DMSO-d₆ 300 MHz) 7.92 (d,
20 J=8.7 Hz, 2H), 7.66 (d, J=8.7 Hz, 2H), 7.49 (s, 2H), 7.46
(dd, J=2.7, 9.6 Hz, 1H), 7.11 (dt, J=2.7, 8.7 Hz, 1H), 6.77
(dd, J=5.7, 8.7 Hz, 1H), 5.72 (t, J=5.7 Hz, 1H), 4.49 (d,
J= 5.7 Hz, 2H), 4.04 (s, 2H). Anal. calc'd. for
C₁₇H₁₄N₃S₂O₃F: C, 52.16; H, 3.61; N, 10.73. Found: C,
25 52.42; H, 3.69; N, 10.49.

The following compounds were obtained according
to procedures similar to that exemplified above and
in the General Synthetic Schemes.

- 37) 4-[7-chloro-1,5-dihydro-3-trifluoromethyl-thieno[3',2':4,5]thiopyrano-s-oxide[3,2-c]pyrazol-1-yl]benzenesulfonamide: mp = 185°C (dec).
- 5 38) 4-[3-(difluoromethyl)-1,5-dihydro-6-fluoro-7-methoxy-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide: mp = 256-258°C.
- 10 39) 4-[1,5-dihydro-7-fluoro-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide: mp = 240-242°C.
- 15 40) [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-fluoro-[2]benzothiopyrano[4,3-c]pyrazol-3-yl]carboxamide: mp = 297-298°C.
- 20 41) 4-[1,5-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-[2]benzothiopyrano-S-oxide[4,3-c]pyrazol-1-yl]benzenesulfonamide: mp = >300°C.
- 25 42) methyl [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-fluoro-[2]benzothiopyrano[4,3-c]pyrazol-3-yl]carboxylate: mp = 241-244°C.
- 25 43) 4-[4,6-dihydro-7-fluoro-8-methoxy-3-(trifluoromethyl)-[2]benzothiepine[5,4-c]pyrazol-1-yl]benzenesulfonamide: mp = 133-138°C.

BIOLOGICAL EVALUATION

30

Rat Carrageenan Foot Pad Edema Test

- The carrageenan foot edema test was performed with materials, reagents and procedures essentially as described by Winter, et al., (Proc. Soc. Exp. Biol. Med., 111, 544 (1962)). Male Sprague-Dawley rats were selected in each group so that the average body weight was as close as possible. Rats were

fasted with free access to water for over sixteen hours prior to the test. The rats were dosed orally (1 mL) with compounds suspended in vehicle containing 0.5% methylcellulose and 0.025% surfactant, or with
5 vehicle alone. One hour later a subplantar injection of 0.1 mL of 1% solution of carrageenan/sterile 0.9% saline was administered and the volume of the injected foot was measured with a displacement plethysmometer connected to a pressure transducer
10 with a digital indicator. Three hours after the injection of the carrageenan, the volume of the foot was again measured. The average foot swelling in a group of drug-treated animals was compared with that of a group of placebo-treated animals and the
15 percentage inhibition of edema was determined (Otterness and Bliven, *Laboratory Models for Testing NSAIDs, in Non-steroidal Anti-Inflammatory Drugs*, (J. Lombardino, ed. 1985)). The % inhibition shows the % decrease from control paw volume
20 determined in this procedure and the data for selected compounds in this invention are summarized in Table XVII.

Rat Carrageenan-induced Analgesia Test

25 The analgesia test using rat carrageenan was performed with materials, reagents and procedures essentially as described by Hargreaves, et al., (Pain, 32, 77 (1988)). Male Sprague-Dawley rats were treated as previously described for the Carrageenan
30 Foot Pad Edema test. Three hours after the injection of the carrageenan, the rats were placed in a special plexiglass container with a transparent floor having a high intensity lamp as a radiant heat source, positionable under the floor. After an initial twenty
35 minute period, thermal stimulation was begun on either the injected foot or on the contralateral uninjected foot. A photoelectric cell turned off the

lamp and timer when light was interrupted by paw withdrawal. The time until the rat withdraws its foot was then measured. The withdrawal latency in seconds was determined for the control and drug-treated groups, and percent inhibition of the hyperalgesic foot withdrawal determined. Results are shown in Table XVII.

TABLE XVII.

10

	RAT PAW EDEMA % Inhibition ¹	ANALGESIA % Inhibition ¹
Example		
15 1	43	35
2	28	29
21	26	14
1- @ 30 mg/kg body weight		

20 Evaluation of COX-1 and COX-2 activity in vitro

The compounds of this invention exhibited inhibition in vitro of COX-2. The COX-2 inhibition activity of the compounds of this invention illustrated in the Examples was determined by the following methods.

25 a. Preparation of recombinant COX baculoviruses

A 2.0 kb fragment containing the coding region of either human or murine COX-1 or human or murine COX-2 was cloned into a BamH1 site of the baculovirus transfer vector pVL1393 (Invitrogen) to generate the baculovirus transfer vectors for COX-1 and COX-2 in a manner similar to the method of D.R. O'Reilly et al (*Baculovirus Expression Vectors: A Laboratory Manual* (1992)). Recombinant baculoviruses were isolated by transfecting 4 µg of baculovirus transfer vector DNA into SF9 insect cells (2x10e8) along with 200 ng of

- linearized baculovirus plasmid DNA by the calcium phosphate method. See M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agric. Exp. Station
- 5 Bull. 1555 (1987). Recombinant viruses were purified by three rounds of plaque purification and high titer (10^7 - 10^8 pfu/ml) stocks of virus were prepared. For large scale production, SF9 insect cells were infected in 10 liter fermentors (0.5×10^6 /ml) with
- 10 the recombinant baculovirus stock such that the multiplicity of infection was 0.1. After 72 hours the cells were centrifuged and the cell pellet homogenized in Tris/Sucrose (50 mM: 25%, pH 8.0) containing 1% 3-[(3-cholamidopropyl)dimethylammonio]-
- 15 1-propanesulfonate (CHAPS). The homogenate was centrifuged at $10,000 \times g$ for 30 minutes, and the resultant supernatant was stored at -80°C before being assayed for COX activity.
- 20 b. Assay for COX-1 and COX-2 activity:
COX activity was assayed as PGE_2 formed/ μg protein/time using an ELISA to detect the prostaglandin released. CHAPS-solubilized insect cell membranes containing the appropriate COX enzyme were
- 25 incubated in a potassium phosphate buffer (50 mM, pH 8.0) containing epinephrine, phenol, and heme with the addition of arachidonic acid ($10 \mu\text{M}$). Compounds were pre-incubated with the enzyme for 10-20 minutes prior to the addition of arachidonic acid. Any
- 30 reaction between the arachidonic acid and the enzyme was stopped after ten minutes at 37°C /room temperature by transferring $40 \mu\text{l}$ of reaction mix into $160 \mu\text{l}$ ELISA buffer and $25 \mu\text{M}$ indomethacin. The PGE_2 formed was measured by standard ELISA technology
- 35 (Cayman Chemical). Results are shown in Table XVIII.

170

TABLE XVIII.

		Human COX-2	Human COX-1
Example		IC ₅₀ μ M	IC ₅₀ μ M
5	1	<0.1	2.7
	2	<0.1	>100
	3	1.3	7.1
	4	0.5	
	5	4.8	24.4
10	6	23.5	11.1
	7	0.5	9.7
	8	0.5	
	9	48.2	>100
	10	<0.1	6.2
15	11	52	>100
	12	95	>100
	13	<0.1	<0.1
	14	0.2	4.6
	15	<0.1	0.3
20	16	<0.1	0.4
	17	<0.1	0.8
	18	<0.1	3.7
	19	<0.1	0.3
	20	<0.1	<0.1
25	21	<0.1	8.7
	22	3.5	51
	24	<0.1	1.2
	25	<0.1	1.5
	26	<0.1	0.2
30	27	59.9	>100
	28	<0.1	>100
	29	0.3	>100
	30	<0.1	10.7
	31	<0.1	0.2
35	32	0.2	8.9
	33	<0.1	<0.1
	34	4.2	>100

TABLE XVIII. (cont.)

Example	Human COX-2	Human COX-1
	IC ₅₀ μ M	IC ₅₀ μ M
5	35	0.1
	36	0.8
	37	1.7
	38	45.6
	39	91.1
10	40	<0.1
	41	>100
	42	0.3
	43	4.7
	44	>100

15

Biological paradigms for testing the cytokine-inhibiting activity of these compounds are found in WO95/13067, published 18 May 1995.

Also embraced within this invention is a class of pharmaceutical compositions comprising the active compounds of this combination therapy in association with one or more non-toxic, pharmaceutically-acceptable carriers and/or diluents and/or adjuvants (collectively referred to herein as "carrier" materials) and, if desired, other active ingredients. The active compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The active compounds and composition may, for example, be administered orally, intravascularly, intraperitoneally, subcutaneously, intramuscularly or topically.

For oral administration, the pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of

a dosage unit containing a particular amount of the active ingredient. Examples of such dosage units are tablets or capsules. The active ingredient may also be administered by injection as a composition wherein, for
5 example, saline, dextrose or water may be used as a suitable carrier.

The amount of therapeutically active compounds that are administered and the dosage regimen for treating a disease condition with the compounds and/or
10 compositions of this invention depends on a variety of factors, including the age, weight, sex and medical condition of the subject, the severity of the disease, the route and frequency of administration, and the particular compound employed, and thus may vary widely.
15 The pharmaceutical compositions may contain active ingredients in the range of about 0.1 to 2000 mg, preferably in the range of about 0.5 to 500 mg and most preferably between about 1 and 100 mg. A daily dose of about 0.01 to 100 mg/kg body weight, preferably between
20 about 0.5 and about 20 mg/kg body weight and most preferably between about 0.1 to 10 mg/kg body weight, may be appropriate. The daily dose can be administered in one to four doses per day.

In the case of psoriasis and other skin
25 conditions, it may be preferable to apply a topical preparation of compounds of this invention to the affected area two to four times a day.

For inflammations of the eye or other external tissues, e.g., mouth and skin, the formulations are
30 preferably applied as a topical ointment or cream, or as a suppository, containing the active ingredients in a total amount of, for example, 0.075 to 30% w/w, preferably 0.2 to 20% w/w and most preferably 0.4 to 15% w/w. When formulated in an ointment, the active
35 ingredients may be employed with either paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an

oil-in-water cream base. If desired, the aqueous phase of the cream base may include, for example at least 30% w/w of a polyhydric alcohol such as propylene glycol, butane-1,3-diol, mannitol, sorbitol, glycerol, polyethylene glycol and mixtures thereof. The topical formulation may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. Examples of such dermal penetration enhancers include dimethylsulfoxide and related analogs. The compounds of this invention can also be administered by a transdermal device. Preferably topical administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. In either case, the active agent is delivered continuously from the reservoir or microcapsules through a membrane into the active agent permeable adhesive, which is in contact with the skin or mucosa of the recipient. If the active agent is absorbed through the skin, a controlled and predetermined flow of the active agent is administered to the recipient. In the case of microcapsules, the encapsulating agent may also function as the membrane.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier, it may comprise a mixture of at least one emulsifier with a fat or an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make-up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations. Emulsifiers and emulsion stabilizers

suitable for use in the formulation of the present invention include Tween 60, Span 80, cetostearyl alcohol, myristyl alcohol, glyceryl monostearate, and sodium lauryl sulfate, among others.

- 5 The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties, since the solubility of the active compound in most oils likely to be used in pharmaceutical emulsion formulations is very low. Thus, the cream
- 10 should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol
- 15 diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters may be used. These may be used alone or in combination depending on the properties required.
- 20 Alternatively, high melting point lipids such as white soft paraffin and/or liquid paraffin or other mineral oils can be used.

- Formulations suitable for topical administration to the eye also include eye drops wherein the active
- 25 ingredients are dissolved or suspended in suitable carrier, especially an aqueous solvent for the active ingredients. The antiinflammatory active ingredients are preferably present in such formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10%
- 30 and particularly about 1.5% w/w.


- For therapeutic purposes, the active compounds of this combination invention are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered *per os*, the
- 35 compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate,

magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations.

What is claimed is:

1. A compound of Formula I



5

wherein A is $-(CH_2)_m-X-(CH_2)_n-$;

wherein X is selected from $S(O)_p$, O and NR^3 ;

wherein m is 0 to 3, inclusive;

10 wherein n is 0 to 3, inclusive;

wherein p is 0 to 2, inclusive;

wherein B is selected from aryl and heteroaryl;

wherein R^1 is selected from hydrido, halo,

haloalkyl, cyano, nitro, formyl, alkoxycarbonyl,

15 carboxyl, carboxyalkyl, alkoxycarbonylalkyl, amidino,

cyanoamidino, aminocarbonyl, alkoxy, alkoxyalkyl,

aminocarbonylalkyl, N-alkylaminocarbonyl, N-

arylamino carbonyl, N,N-dialkylaminocarbonyl, N-alkyl-

N-arylamino carbonyl, alkylcarbonyl,

20 alkylcarbonylalkyl, hydroxyalkyl, alkylthio,

alkylsulfinyl, alkylsulfonyl, alkylthioalkyl,

alkylsulfinylalkyl, alkylsulfonylalkyl, N-

alkylaminosulfonyl, N-arylamino sulfonyl, arylsulfonyl,

N,N-dialkylaminosulfonyl, N-alkyl-N-arylamino sulfonyl

25 and heterocyclic;

wherein R^2 is selected from aryl and heteroaryl,

wherein R^2 is optionally substituted at a

substitutable position with one or more radicals

selected from alkylsulfonyl, aminosulfonyl, halo,

30 alkyl, alkoxy, hydroxyl and haloalkyl;

wherein R^3 is selected from hydrido and alkyl;

and

wherein R^4 is one or more radicals selected from

hydrido, halo, alkylthio, alkylsulfinyl, alkyl,

alkylsulfonyl, cyano, carboxyl, alkoxycarbonyl, amido, N-alkylaminocarbonyl, N-arylaminocarbonyl, N,N-dialkylaminocarbonyl, N-alkyl-N-arylaminocarbonyl, haloalkyl, hydroxyl, alkoxy, hydroxyalkyl, haloalkoxy, 5 aminosulfonyl, N-alkylaminosulfonyl, amino, N-alkylamino, N,N-dialkylamino, heterocyclic, nitro and acylamino;

provided either R⁴ is aminosulfonyl or alkylsulfonyl, or R² is substituted with aminosulfonyl 10 or alkylsulfonyl;
or a pharmaceutically-acceptable salt thereof.

2. Compound of Claim 1 wherein A is $-(CH_2)_m-X-(CH_2)_n-$; wherein X is selected from S(O)_p, O and NR³; 15 wherein m is 0 to 3, inclusive; wherein n is 0 to 3, inclusive; wherein p is 0 to 2, inclusive; wherein B is selected from phenyl, naphthyl and five and six membered heteroaryl; wherein R¹ is selected from halo, lower haloalkyl, cyano, nitro, formyl, lower 20 alkoxycarbonyl, lower carboxyalkyl, lower alkoxycarbonylalkyl, amidino, cyanoamidino, lower alkoxy, lower alkoxyalkyl, aminocarbonyl, lower aminocarbonylalkyl, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl, 25 lower N-alkyl-N-phenylaminocarbonyl, lower alkylcarbonyl, lower alkylcarbonylalkyl, lower hydroxyalkyl, lower alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, lower alkylthioalkyl, lower alkylsulfinylalkyl, lower alkylsulfonylalkyl, lower N- 30 alkylaminosulfonyl, N-phenylaminosulfonyl, phenylsulfonyl, lower N,N-dialkylaminosulfonyl, lower N-alkyl-N-phenylaminosulfonyl and five-seven membered heterocyclic; wherein R² is selected from phenyl and five or six membered heteroaryl, wherein R² is 35 optionally substituted at a substitutable position with one or more radicals selected from lower alkylsulfonyl, aminosulfonyl, halo, lower alkyl, lower

alkoxy, hydroxyl and lower haloalkyl; wherein R^3 is selected from hydrido and lower alkyl; and wherein R^4 is one or more radicals selected from hydrido, halo, lower alkylthio, lower alkylsulfinyl, lower alkyl, lower alkylsulfonyl, cyano, carboxyl, lower alkoxycarbonyl, aminocarbonyl, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl, lower N-alkyl-N-phenylaminocarbonyl, lower haloalkyl, hydroxyl, lower alkoxy, lower hydroxyalkyl, lower haloalkoxy, aminosulfonyl, lower N-alkylaminosulfonyl, amino, lower N-alkylamino, lower N,N-dialkylamino, five-seven membered heterocyclic, nitro and acylamino; or a pharmaceutically-acceptable salt thereof.

15

3. Compound of Claim 2 wherein A is $-(CH_2)_m-X-(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0, 1 or 2; wherein n is 0, 1 or 2; wherein p is 0, 1 or 2; wherein B is selected from phenyl and five and six membered heteroaryl; wherein R^1 is selected from halo, lower haloalkyl, cyano, formyl, lower alkoxycarbonyl, aminocarbonyl, lower alkoxycarbonylalkyl, lower alkoxy, lower alkoxyalkyl, lower aminocarbonylalkyl, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl, lower N-alkyl-N-phenylaminocarbonyl and lower hydroxyalkyl; wherein R^2 is phenyl substituted at a substitutable position with a radical selected from lower alkylsulfonyl and aminosulfonyl; and wherein R^4 is one or more radicals selected from hydrido, halo, lower alkylthio, lower alkylsulfinyl, lower alkyl, lower alkylsulfonyl, cyano, carboxyl, lower alkoxycarbonyl, aminocarbonyl, lower N-alkylaminocarbonyl, lower N,N-dialkylaminocarbonyl, lower N-alkyl-N-phenylaminocarbonyl, lower haloalkyl, hydroxyl, lower alkoxy, lower hydroxyalkyl, amino, lower N-alkylamino,

lower N,N-dialkylamino, lower haloalkoxy and nitro; or a pharmaceutically-acceptable salt thereof.

4. Compound of Claim 3 wherein A is $-(CH_2)_m-X-$
5 $(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from phenyl and five and six membered heteroaryl; wherein R^1 is selected from lower haloalkyl, lower hydroxyalkyl, cyano, formyl, lower
10 alkoxy, carbonyl, lower alkoxy, lower N-alkylaminocarbonyl, N-phenylaminocarbonyl, lower N,N-dialkylaminocarbonyl and lower N-alkyl-N-phenylaminocarbonyl; wherein R^2 is phenyl substituted at a substitutable position with a radical selected
15 from lower alkylsulfonyl and aminosulfonyl; and wherein R^4 is one or more radicals selected from hydrido, halo, lower alkylthio, lower alkylsulfinyl, lower alkyl, cyano, carboxyl, lower alkoxy, carbonyl, lower alkoxy, amino, lower N-alkylamino, lower N,N-dialkylamino, lower hydroxyalkyl and lower haloalkoxy;
20 or a pharmaceutically-acceptable salt thereof.

5. Compound of Claim 4 wherein A is $-(CH_2)_m-X-$
25 $(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0 or 1; wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from phenyl, thienyl, pyridyl, furyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, pyridazinyl,
30 pyrimidinyl, pyrazinyl, triazinyl, thiaimidazolyl, oxoimidazolyl, azaoxazolyl, azathiazolyl and pyrrolyl; wherein R^1 is selected from fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl,
35 heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, hydroxymethyl,

- hydroxyethyl, cyano, formyl, carboxyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tert-butoxycarbonyl, propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl, aminocarbonyl,
- 5 methoxy, ethoxy, propoxy, n-butoxy, N-methylaminocarbonyl, N-phenylaminocarbonyl, N,N-dimethylaminocarbonyl, N-methyl-N-phenylaminocarbonyl and methylcarbonyl; wherein R² is phenyl substituted at a substitutable position with a radical selected
- 10 from methylsulfonyl and aminosulfonyl; wherein R⁴ is optionally substituted with one or more radicals selected from hydrido, fluoro, chloro, bromo, methylthio, ethylthio, isopropylthio, tert-butylthio, isobutylthio, hexylthio, methylsulfinyl,
- 15 ethylsulfinyl, isopropylsulfinyl, tert-butylsulfinyl, isobutylsulfinyl, hexylsulfinyl, methyl, ethyl, isopropyl, tert-butyl, isobutyl, hexyl, cyano, carboxyl, methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl, tert-butoxycarbonyl,
- 20 propoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, pentoxycarbonyl, aminocarbonyl, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl,
- 25 dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, hydroxyl, methoxy, methylenedioxy, ethoxy, propoxy, n-butoxy, hydroxymethyl and trifluoromethoxy; or a pharmaceutically-acceptable salt thereof.
- 30

6. Compound of Claim 5 selected from compounds, and their pharmaceutically acceptable salts, of the group consisting of

- 35 6-fluoro-7-methoxy-1-[(4-methylsulfonyl)phenyl]-3-(trifluoromethyl)-1H-[1]benzothieno[3,2-c]pyrazole;

- 4-[6-fluoro-7-methoxy-3-(trifluoromethyl)-1H-[1]benzothieno[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 5 4-[6-methoxy-3-(trifluoromethyl)-1H-[1]benzothieno[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[7-fluoro-3-(trifluoromethyl)-1H-[1]benzothieno[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[7-chloro-3-(trifluoromethyl)-1H-[1]benzothieno[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 10 4-[7-methyl-3-(trifluoromethyl)-1H-[1]benzothieno[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[3-(trifluoromethyl)-1H-[1]benzothieno[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 15 6-fluoro-7-methoxy-1-[(4-methylsulfonyl)phenyl]-3-(trifluoromethyl)-1H-[1]benzofuro[3,2-c]pyrazole;
- 4-[6-fluoro-7-methoxy-3-(trifluoromethyl)-1H-[1]benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[6-methoxy-3-(trifluoromethyl)-1H-[1]benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 20 4-[7-fluoro-3-(trifluoromethyl)-1H-[1]benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[7-chloro-3-(trifluoromethyl)-1H-[1]benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 25 4-[7-methyl-3-(trifluoromethyl)-1H-[1]benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[3-(trifluoromethyl)-1H-benzofuro[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 1,5-dihydro-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)-[2]benzopyrano[4,3-c]pyrazole;
- 30 1,5-dihydro-7-methyl-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)-[2]benzopyrano[4,3-c]pyrazole;
- 1,5-dihydro-6-fluoro-7-methoxy-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazole;
- 35

- 4-[1,5-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,5-dihydro-3-(trifluoromethyl)-[2]benzopyrano[4,3-
c]pyrazol-1-yl]benzenesulfonamide;
- 5 4-[1,5-dihydro-7-fluoro-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[6-chloro-1,5-dihydro-7-methoxy-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 10 4-[7-chloro-1,5-dihydro-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 15 4-[1,5-dihydro-7-methoxy-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[6,7-difluoro-1,5-dihydro-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 20 4-[1,5-dihydro-3-(trifluoromethyl)-[2]benzopyrano[4,3-
c]pyrazol-1-yl]benzenesulfonamide;
- 4-[3-(difluoromethyl)-1,5-dihydro-6-fluoro-7-methoxy-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 25 1,4-dihydro-1-[4-(methylsulfonyl)phenyl]-3-
(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazole;
- 1,4-dihydro-7-methyl-1-[4-(methylsulfonyl)phenyl]-3-
(trifluoromethyl)-[1]benzopyrano[4,3-c]pyrazole;
- 30 1,4-dihydro-6-fluoro-7-methoxy-1-[4-
(methylsulfonyl)phenyl]-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazole;
- 4-[1,4-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-
[1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 35 4-[1,4-dihydro-3-(trifluoromethyl)-[1]benzopyrano[4,3-
c]pyrazol-1-yl]benzenesulfonamide;

- 4-[1,4-dihydro-7-fluoro-3-(trifluoromethyl)-
[1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[6-chloro-1,4-dihydro-7-methoxy-3-(trifluoromethyl)-
5 [1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[7-chloro-1,4-dihydro-3-(trifluoromethyl)-
[1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 10 4-[1,4-dihydro-7-methoxy-3-(trifluoromethyl)-
[1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,4-dihydro-7-methyl-3-(trifluoromethyl)-
[1]benzopyrano[4,3-c]pyrazol-1-
15 yl]benzenesulfonamide;
- 4-[6,7-difluoro-1,4-dihydro-3-(trifluoromethyl)-
[1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[3-(difluoromethyl)-1,4-dihydro-6-fluoro-7-methoxy-
20 [1]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,4-dihydro-3-(trifluoromethyl)-[1]
benzopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[3-(difluoromethyl)-1,5-dihydro-7-methyl-
25 [2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,5-dihydro-7,8,9-trimethoxy-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 30 4-[1,5-dihydro-7-methyl-3-(trifluoromethyl)-
[2]benzopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- methyl [1-[4-(aminosulfonyl)phenyl]-1,4-dihydro-
[1]benzopyrano[4,3-c]pyrazol-3-yl]carboxylate;
- 35 methyl [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-
fluoro-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-
c]pyrazol-3-yl]carboxylate;

- meth] [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-fluoro-[2]benzothiopyrano[4,3-c]pyrazol-3-yl]carboxylate;
- 4-[7-chloro-3-(difluoromethyl)-1,5-dihydro-
- 5 [2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,5-dihydro-3-(trifluoromethyl)-[1,3]dioxolo[6,7][2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 10 4-[7-fluoro-1,5-dihydro-3-(hydroxymethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[3-(difluoromethyl)-1,5-dihydro-7-methyl-
- 15 [2]benzenethiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[3-cyano-7-fluoro-1,5-dihydro-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]-N-[(dimethylamino)methylene]benzenesulfonamide;
- 4-[3-(difluoromethyl)-7-fluoro-1,5-dihydro-
- 20 [2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[3-cyano-7-fluoro-1,5-dihydro-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 25 4-[6,8-difluoro-1,5-dihydro-7-methoxy-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 7-fluoro-1,5-dihydro-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazole;
- 30 [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-fluoro-[2]benzothiopyrano[4,3-c]pyrazol-3-yl]carboxamide;
- [1-(4-aminosulfonylphenyl)-1,5-dihydro-7-fluoro-3-(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazol-3-yl]carboxamide;
- 35 4-[3-(difluoromethyl)-1,5-dihydro-6-fluoro-7-methoxy-[2]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;

- 4-[1,5-dihydro-7-fluoro-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,5-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-
5 [2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 1,5-dihydro-6-fluoro-7-methoxy-1-[4-
(methylsulfonyl)phenyl]-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazole;
- 10 4-[1,5-dihydro-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[1,5-dihydro-7-methyl-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
15 yl]benzenesulfonamide;
- 1-[1,5-dihydro-4-(methylsulfonyl)phenyl]-7-methyl-3-
(trifluoromethyl)-[2]benzothiopyrano[4,3-c]pyrazole;
- 4-[7-chloro-1,5-dihydro-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
20 yl]benzenesulfonamide;
- 4-[1,5-dihydro-7-methoxy-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[7-chloro-1,5-dihydro-3-trifluoromethyl-
25 thieno[3',2':4,5]thiopyrano[3,2-c]pyrazol-1-
yl]benzenesulfonamide;
- 4-[7-chloro-1,5-dihydro-3-trifluoromethyl-
thieno[3',2':4,5]thiopyrano-S-oxide[3,2-c]pyrazol-1-
yl]benzenesulfonamide;
- 30 4-[1,5-dihydro-3-
(trifluoromethyl)furano[2',3':4,5]thiopyrano[3,2-
c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,5-dihydro-3-
(trifluoromethyl)oxazolo[4',5':4,5]thiopyrano[3,2-
35 c]pyrazol-1-yl]benzenesulfonamide;

- 4-[1,5-dihydro-3-(trifluoromethyl)thiazolo[5',4':4,5]thiopyrano[3,2-c]pyrazol-1-yl]benzenesulfonamide;
- 1,5-dihydro-1-(4-methoxyphenyl)-3-
- 5 (trifluoromethyl)thieno[3',2':4,5]thiopyrano[3,2-c]pyrazole-7-sulfonamide;
- 4-[1,5-dihydro-3-(trifluoromethyl)pyrazolo[3',4':5,6]thiopyrano[3,4-b]pyridin-1-yl]benzenesulfonamide;
- 10 4-[1,5-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-[2]benzothiopyrano-S-oxide[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,4-dihydro-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-
- 15 yl]benzenesulfonamide;
- methyl [1-[4-(aminosulfonyl)phenyl]-1,4-dihydro-[1]benzothiopyrano[4,3-c]pyrazol-3-carboxylate;
- 4-[6,7-dichloro-1,4-dihydro-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-
- 20 yl]benzenesulfonamide;
- 4-[1,4-dihydro-7-fluoro-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,4-dihydro-6-isopropyl-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-
- 25 yl]benzenesulfonamide;
- 4-[1,4-dihydro-7,8-dimethoxy-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 30 4-[1,4-dihydro-7-methoxy-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-yl]benzenesulfonamide;
- 4-[1,4-dihydro-7-methyl-3-(trifluoromethyl)-[1]benzothiopyrano[4,3-c]pyrazol-1-
- 35 yl]benzenesulfonamide;


- 4-[6-chloro-1,4-dihydro-7-methoxy-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
4-[1,4-dihydro-6-fluoro-7-methoxy-3-(trifluoromethyl)-
5 [1]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
4-[7-chloro-1,4-dihydro-3-(trifluoromethyl)-
[1]benzothiopyrano[4,3-c]pyrazol-1-
yl]benzenesulfonamide;
10 4-[4,6-dihydro-7-fluoro-8-methoxy-3-(trifluoromethyl)-
[1]benzothiepino[5,4-c]pyrazol-1-
yl]benzenesulfonamide; and
1-(4-aminosulfonylphenyl)-1,4-
dihydro[1]benzothiopyrano[4,3-c]pyrazole-3-
15 yl]carbonitrile.

7. Compound of Claim 5 which is 4-[1,5-dihydro-6-
fluoro-7-methoxy-3-(trifluoromethyl)-
[2]benzothiopyrano[4,3-c]pyrazol-1-
20 yl]benzenesulfonamide, or a pharmaceutically-
acceptable salt thereof.

8. Compound of Claim 5 which is 1-[6-fluoro-7-
methoxy-4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)-
25 1,5-dihydro-[2]benzothiopyrano[4,3-c]pyrazole, or a
pharmaceutically-acceptable salt thereof.

9. Compound of Claim 5 which is 4-[7-chloro-1,5-
dihydro-3-trifluoromethyl-[2]thienothiopyrano[4,3-
30 c]pyrazol-1-yl]benzenesulfonamide, or a
pharmaceutically-acceptable salt thereof.

10. A compound of Formula II



- 5 wherein A is $-(CH_2)_m-X-(CH_2)_n-$;
 wherein X is $S(O)_p$ or O;
 wherein m is 0 or 1;
 wherein n is 0 or 1;
 wherein p is 0 or 1;
 10 wherein B is selected from aryl and heteroaryl;
 wherein R^1 is selected from haloalkyl,
 hydroxyalkyl, aminocarbonyl, alkoxy carbonyl and cyano;
 wherein R^4 is one or more radicals selected from
 hydrido, halo, alkyl and alkoxy; and
 15 wherein R^5 is selected from alkyl and amino;
 or a pharmaceutically-acceptable salt thereof.

11. Compound of Claim 10 wherein A is $-(CH_2)_m-X-$
 $(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0 or 1;
 20 wherein n is 0 or 1; wherein p is 0 or 1; wherein B is
 selected from phenyl and five membered heteroaryl;
 wherein R^1 is selected from lower haloalkyl, lower
 hydroxyalkyl, aminocarbonyl, lower alkoxy carbonyl and
 cyano; wherein R^4 is one or more radicals selected
 25 from hydrido, halo, lower alkyl and lower alkoxy; and
 wherein R^5 is selected from lower alkyl and amino; or
 a pharmaceutically-acceptable salt thereof.

12. Compound of Claim 11 wherein A is $-(CH_2)_m-X-$
 30 $(CH_2)_n-$; wherein X is $S(O)_p$ or O; wherein m is 0 or 1;

wherein n is 0 or 1; wherein p is 0 or 1; wherein B is selected from phenyl, thienyl, furyl and pyrrolyl; wherein R¹ is selected from fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, hydroxymethyl, hydroxyethyl, aminocarbonyl, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and cyano; wherein R⁴ is one or more radicals selected from hydrido, fluoro, chloro, bromo, methyl, ethyl, isopropyl, tert-butyl, isobutyl, hexyl, methoxy, methylenedioxy, ethoxy, propoxy, n-butoxy and tert-butoxy; and wherein R⁵ is selected from methyl and amino; or a pharmaceutically-acceptable salt thereof.

13. A pharmaceutical composition comprising a therapeutically-effective amount of a compound, said compound selected from a family of compounds of Claim 1, 2, 3, 4, 5, 6, 7, 8 or 9; or a pharmaceutically-acceptable salt thereof.

14. A method of treating inflammation or an inflammation-associated disorder in a subject, said method comprising administering to the subject having or susceptible to said inflammation or inflammation-associated disorder, a therapeutically-effective amount of a compound of Claim 1, 2, 3, 4, 5, 6, 7, 8 or 9; or a pharmaceutically-acceptable salt thereof.

15. The method of Claim 14 for use in treatment of inflammation.

16. The method of Claim 14 for use in treatment of an inflammation-associated disorder.

17. The method of Claim 16 wherein the inflammation-associated disorder is arthritis.

18. The method of Claim 16 wherein the
5 inflammation-associated disorder is pain.

19. The method of Claim 16 wherein the inflammation-associated disorder is fever.

INTERNATIONAL SEARCH REPORT

 Interno J Application No
 PCT/US 95/11403

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 6 C07D491/048 A61K31/415 C07D491/052 C07D495/04 C07D495/14
 C07D513/14 //(C07D491/048, 307:00, 231:00), (C07D495/04, 333:00,
 231:00), (C07D491/052, 311:00, 231:00), (C07D495/04, 335:00, 231:00),

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 347 773 (FARMITALIA) 27 December 1989 cited in the application see claims 1,6 ---	1,10,13
A	EP,A,0 203 679 (E. I. DU PONT) 3 December 1986 see page 193 - page 195; claim 1 -----	1,10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

27 December 1995

Date of mailing of the international search report

9.01.96

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Voyiazoglou, D

INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/US 95/11403

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 (C07D495/04, 337:00, 231:00), (C07D495/14, 335:00, 231:00, 221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

27 December 1995

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Voyiazoglou, D

INTERNATIONAL SEARCH REPORT

Int. l. application No.

PCT/US95/11403

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 14-19 are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 95/11403

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0347773	27-12-89	AU-B- 634966	11-03-93
		AU-B- 3769289	12-01-90
		CN-B- 1022322	06-10-93
		WO-A- 8912630	28-12-89
		EP-A- 0422051	17-04-91
		ES-T- 2054930	16-08-94
		IE-B- 63189	22-03-95
		IL-A- 90552	26-08-94
		JP-T- 3505205	14-11-91
		PT-B- 90905	30-12-94
		US-A- 5424308	13-06-95
EP-A-0203679	03-12-86	US-A- 4678499	07-07-87
		CA-A- 1230334	15-12-87
		US-A- 4741765	03-05-88

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.